

## Math 1553 Worksheet §§3.5-4.3

### Solutions

1. True or false. Answer true if the statement is *always* true. Otherwise, answer false. If your answer is false, either give an example that shows it is false or (in the case of an incorrect formula) state the correct formula.
  - a) If  $A$  and  $B$  are  $n \times n$  matrices and both are invertible, then the inverse of  $AB$  is  $A^{-1}B^{-1}$ .
  - b) If  $A$  and  $B$  are invertible  $n \times n$  matrices, then  $A + B$  is invertible and
 
$$(A + B)^{-1} = A^{-1} + B^{-1}.$$
  - c) Suppose  $A$  is an  $n \times n$  matrix and every vector in  $\mathbf{R}^n$  can be written as a linear combination of the columns of  $A$ . Then  $A$  must be invertible.
  - d) If  $\det(A) = 1$  and  $c$  is a scalar, then  $\det(cA) = c \det(A)$ .

### Solution.

- a) False.  $(AB)^{-1} = B^{-1}A^{-1}$ .
- b) False.  $A + B$  might not be invertible in the first place. For example, if  $A = I_2$  and  $B = -I_2$  then  $A + B = 0$  which is not invertible. Even in the case when  $A + B$  is invertible, it still might not be true that  $(A + B)^{-1} = A^{-1} + B^{-1}$ . For example,  $(I_2 + I_2)^{-1} = (2I_2)^{-1} = \frac{1}{2}I_2$ , whereas  $(I_2)^{-1} + (I_2)^{-1} = I_2 + I_2 = 2I_2$ .
- c) True. If the columns of  $A$  span  $\mathbf{R}^n$ , then  $A$  is invertible by the Invertible Matrix Theorem. We can also see this directly without quoting the IMT:  
 If the columns of  $A$  span  $\mathbf{R}^n$ , then  $A$  has  $n$  pivots, so  $A$  has a pivot in each row and column, hence its matrix transformation  $T(x) = Ax$  is one-to-one and onto and thus invertible. Therefore,  $A$  is invertible.
- d) False. By the properties of the determinant, scaling one row by  $c$  multiplies the determinant by  $c$ . When we take  $cA$  for an  $n \times n$  matrix  $A$ , we are multiplying *each* row by  $c$ . This multiplies the determinant by  $c$  a total of  $n$  times. Thus, if  $A$  is  $n \times n$  and  $\det(A) = 1$ , then

$$\det(cA) = c^n \det(A) = c^n(1) = c^n.$$

2. Let  $A = \begin{pmatrix} 7 & 1 & 4 & 1 \\ -1 & 0 & 0 & 6 \\ 9 & 0 & 2 & 3 \\ 0 & 0 & 0 & -1 \end{pmatrix}$

- a) Compute  $\det(A)$ .
- b) Compute  $\det(A^{-1})$  without doing any more work.
- c) Compute  $\det((A^T)^5)$  without doing any more work.
- d) Find the volume of the parallelepiped formed by the columns of  $A$ .

**Solution.**

a) The second column has three zeros, so we expand by cofactors:

$$\det \begin{pmatrix} 7 & 1 & 4 & 1 \\ -1 & 0 & 0 & 6 \\ 9 & 0 & 2 & 3 \\ 0 & 0 & 0 & -1 \end{pmatrix} = -\det \begin{pmatrix} -1 & 0 & 6 \\ 9 & 2 & 3 \\ 0 & 0 & -1 \end{pmatrix}$$

Now we expand the second column by cofactors again:

$$\dots = -2 \det \begin{pmatrix} -1 & 6 \\ 0 & -1 \end{pmatrix} = (-2)(-1)(-1) = -2.$$

b) From our notes, we know  $\det(A^{-1}) = \frac{1}{\det(A)} = -\frac{1}{2}$ .

c)  $\det(A^T) = \det(A) = -2$ . By the multiplicative property of determinants, if  $B$  is any  $n \times n$  matrix, then  $\det(B^n) = (\det B)^n$ , so

$$\det((A^T)^5) = (\det A^T)^5 = (-2)^5 = -32.$$

d) Volume of the parallelepiped is  $|\det(A)| = 2$

3. Suppose we have

$$\det \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix} = 5.$$

Compute

$$\det \begin{pmatrix} d-3a & e-3b & f-3c \\ a & b & c \\ 2g & 2h & 2i \end{pmatrix}.$$

**Solution.**

a) Recall that the only operations that affect determinants are "row-swaps" and "row scaling". "Row replacement (through row operation)" does not change the determinant. We have the following row operations:

$$\begin{aligned} \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix} &\xrightarrow{R_1 \leftrightarrow R_2} \begin{pmatrix} d & e & f \\ a & b & c \\ g & h & i \end{pmatrix} \xrightarrow{R_1 = R_1 - 3R_2} \begin{pmatrix} d-3a & e-3b & f-3c \\ a & b & c \\ g & h & i \end{pmatrix} \\ &\xrightarrow{R_3 = 2R_3} \begin{pmatrix} d-3a & e-3b & f-3c \\ a & b & c \\ 2g & 2h & 2i \end{pmatrix}. \end{aligned}$$

Therefore, we have one "row swap" which scales our determinant by  $-1$  and we have a "row multiplied by a scalar" which scales our determinant by the scalar, which in this case is 2. Therefore, we have that

$$\det \begin{pmatrix} d-3a & e-3b & f-3c \\ a & b & c \\ 2g & 2h & 2i \end{pmatrix} = (5)(-1)(2) = -10.$$