

ON SYMPLECTIC FILLINGS

JOHN B. ETNYRE

ABSTRACT. In this note we make several observations concerning symplectic fillings. In particular we show that a (strongly or weakly) semi-fillable contact structure is fillable and any filling embeds as a symplectic domain in a closed symplectic manifold. We also relate properties of the open book decomposition of a contact manifold its possible fillings. These results are also useful in showing the contact Heegaard Floer invariant of a fillable contact structure do not vanish [27] and property P for knots [17].

1. INTRODUCTION

It is important to consider various notion of “symplectic filling” when studying contact and symplectic manifolds (especially in low dimensions). In particular, it is useful to understand when a contact structure on a three manifold has a symplectic filling, what type of filling is it and if there are restrictions on the topology of the filling.

Given a contact three manifold (M, ξ) and symplectic manifold (X, ω) with M a boundary component of X and $\omega|_M > 0$. In particular (X, ω) is a weak (or possibly strong) semi-filling of (M, ξ) . (Throughout this paper all contact structures will be (co)oriented, contact manifolds will be oriented by the contact structures and symplectic manifolds will be oriented by their symplectic structures.) Then we say the boundary component M of X can be *symplectically capped off* if there is a symplectic manifold (X', ω') such that $X' \setminus M = X \cup C$ and $\omega'|_X = \omega$. The symplectic manifold $(C, \omega'|_C)$ is called a symplectic cap for $(M, \xi) \subset (X, \omega)$. (The notation $A \setminus B$ means the union of the metric completions of the components of A minus B .) Many of our results rely on the following theorem.

Theorem 1.1. *If (X, ω) is a (weak or strong) symplectic filling (or semi-filling) of (M, ξ) then the boundary component M of X can be capped off symplectically.*

We have the following immediate corollaries.

Corollary 1.2. *A weakly, respectively strongly, semi-fillable contact structure is fillable.*

Corollary 1.3. *Any symplectic four manifold with weakly or strongly convex boundary components can be embedded as a domain in a closed symplectic manifold.*

The theorem and both corollaries have also been established by Eliashberg in [2]. For *strong* (semi)fillings these results were essentially proven by Gay in [12] and follow from Theorem 1.3 in [10]. Moreover, after writing this paper the author learned that Stipsicz and Ghiggini also know how to prove these results for strong fillings. Moreover, in [28], Stipsicz has and different proof of Lemma 3.1 (below), which is the key ingredient in the proof of Theorem 1.1. In [22], Lisca and Matić proved the corollaries for Stein fillings. Not only do these corollaries illuminate the nature of symplectic and contact geometry in low dimensions they have many

1991 *Mathematics Subject Classification.* Primary 53C15; Secondary 57M50.

Key words and phrases. tight, symplectic filling, convexity.

important applications. Specifically, Kronheimer and Mrowka [17] can use Corollary 1.2 to prove the Property P conjecture for knots and Ozsváth and Szabó [27] can use Corollary 1.3 to show that their Heegaard Floer contact invariant does not vanish for fillable contact structures. Moreover, they can use the corollary to give an alternate proof that if p surgery on a knot K yields a manifold which is (orientation preserving) homeomorphic to $L(p, 1)$ then K is the unknot (this result first appeared in [19] using Seiberg-Witten Floer Homology).

We now turn to the topology of a symplectic filling. A fundamental result of Giroux [13] (see also [15]) says that a contact structure ξ on a three manifold M is always supported by an open book (L, ϕ) . That is, comes from the construction of Thurston and Winkelnkemper [29]; or more explicitly, there is a link L in M that is transverse to ξ such that $M \setminus L$ is fibered by Seifert surfaces for L and the contact structure can be isotoped to be arbitrarily close to the fibers in the fibration (while keeping L transverse). The *monodromy* of the open book ϕ is the monodromy of the fibration $M \setminus L$.

Theorem 1.4. *Suppose the contact structure ξ on a three manifold M is supported by the open book (L, ϕ) and there is a diffeomorphism ψ of the fiber such that $(L, \psi \circ \phi)$ is an open book for S^3 and ψ can be written as a composition of positive Dehn twists. If (X, ω) is a, weak or strong, symplectic semi-filling of (M, ξ) then (X, ω) is a filling of (M, ξ) . Moreover, if M is a rational homology sphere then $H_1(X; \mathbb{Z})$ finite (and trivial if M is an integral homology sphere) and the intersection form on $H_2(X; \mathbb{Z})$ is negative definite.*

Lisca (in [20, 21]) and Ohta-Ono (in [26]) have obtained a similar result under the assumption that M admits a metric of positive scalar curvature (but no assumption on the supporting open book). We note that if ϕ^{-1} can be written as a product of positive Dehn twists then the conditions of the theorem are satisfied (see the proof of Lemma 3.1). If ϕ^{-1} can be so written then ϕ can be written as product of negative Dehn twists. One might expect that this implies the associated contact structure is overtwisted, but this is not necessarily true, only if the open book “destabilizes.” In general it is quite subtle to decide if an open book destabilizes. It is also difficult to make general statements about representing a diffeomorphism in terms of Dehn twists.

Theorem 1.5. *Under the hypothesis of Theorem 1.4 any symplectic filling of (M, ξ) can be embedded as a symplectic domain in a rational surface.*

While these theorems seem to provide potentially quite useful insights into the monodromy of open books we content ourselves here with the following simple applications.

Corollary 1.6. *Let ϕ be any monodromy associated to any tight contact structure coming from perturbing a Reebless foliation into a contact structure as in [8]. Then ϕ cannot be expressed as a composition of negative Dehn twists. Moreover, there is no diffeomorphism ψ such that $\phi \circ \psi$ is the monodromy for an open book of S^3 and ψ is a composition of positive Dehn twists.*

As a specific example any monodromy map associated to any tight contact structure on T^3 cannot be expressed as the composition of negative Dehn twists.

Proof. Such contact structures have semi-fillings with two boundary components. □


Acknowledgments: I thank Yasha Eliashberg for sharing an early version of [2] with me which made me aware of the many interesting applications of symplectic caps. I also gratefully acknowledge support from an NSF CAREER Grant (DMS-0239600).

2. SYMPLECTIC FILLINGS

We recall the various of symplectic fillings of contact manifolds. (For more details, see the survey paper [9], and for more discussion of the notions of fillability see [2].)

A symplectic manifold (X, ω) is said to have *strongly convex boundary* if there is a vector field v defined in the neighborhood of ∂X that points transversely out of X and dilates ω (ie, $\mathcal{L}_v \omega = \omega$). The form $\alpha = (\iota_v \omega)|_{\partial X}$ is a contact form on ∂X . The contact manifold $(M = \partial X, \xi)$ on the boundary is said to be *strongly symplectically fillable* by (X, ω) . If (X, ω) supports a Stein structure then (M, ξ) is said to be *holomorphically fillable* (this is sometimes called Stein fillable). A symplectic manifold (X, ω) is said to have *weakly convex boundary* if ∂X admits a contact structure ξ such that $\omega|_{\xi} > 0$. A contact manifold (M, ξ) is *weakly symplectically fillable* if it is the weakly convex boundary of a symplectic manifold. A contact structure is said to be weakly (or strongly) *semi-fillable* if it is one component of the boundary of a weakly (or strongly) convex symplectic manifold. (Note a Stein manifold always has a connected boundary so there is no notion of holomorphically semi-fillable.) When talking about strong or weak symplectic fillings we will frequently drop the word “symplectic” and just refer to strong or weak fillings.

The following diagram indicates the hierarchy of contact structures.



The proper inclusion of the set of weakly symplectically semi-fillable contact structures into the set of tight contact structures was shown in [11] (for further examples see [23]). The proper inclusion of the set of strongly fillable structures into the set of weakly fillable structures is already seen on T^3 . This is due to Eliashberg [3]. Further results can be found in [1]. The equalities of the semi-fillings and “honest” fillings is the content of Corollary 1.2 and [2]. Though there are strong fillings of a contact structure that are not holomorphic fillings (because there is more than one boundary component [24]) it is not currently known if strongly fillable contact structures are also holomorphically fillable.

If M is a rational homology sphere, a weak symplectic filling can be modified into a strong symplectic filling [26] (the germ of the argument originally appeared in [5], see also [2]). If M has a positive scalar curvature metric, then moreover a semi-filling is automatically a one boundary component filling, and all four notions of symplectic filling become the same (due to Lisca [20] and Ohta-Ono [26]). The proof builds on the work of Kronheimer and Mrowka [18] and relies on Seiberg-Witten theory.

3. PROOF OF THEOREM 1.1

For the proof of Theorem 1.1 we will need the following lemma (see also [28]).

Lemma 3.1. *Under the hypothesis of Theorem 1.1 two handles may be attached to V to form a new symplectic manifold with weakly convex boundary (X', ω') such that $X' \setminus M = X \cup B$ with $\partial B = M' \cup (-M)$ where M' is a homology sphere.*

With this lemma in hand we prove Theorem 1.1

Proof of Theorem 1.1. Given (M, ξ) and (X, ω) as in the theorem, apply the above lemma to obtain (X', ω') . Since M' is a homology sphere the results of Ohta-Ono mentioned in the previous section imply ω' may be isotoped near M' so that (X', ω') is strongly convex along M' . Theorem 1.3 from [10] says that any contact manifold has (infinitely many) strongly concave fillings. Since a strongly convex and strongly concave filling may be glued to form a symplectic manifold [9] we see that (X', ω') may be symplectically capped off. Let C be the cap. Then $C \cup B$ (B from Lemma 3.1) is the symplectic cap for (M, ξ) . \square

We are left to prove the lemma.

Proof of Lemma 3.1. Giroux's result concerning open books and contact structures implies we can assume that (M, ξ) is supported by an open book with connected binding (this can be achieved by "positive Hopf stabilization"). Let ϕ be the monodromy of the open book. Since the mapping class group is generated by right handed Dehn twists about arbitrary loops plus arbitrary Dehn twists parallel to the boundary, we can write ϕ^{-1} as $\phi_0 \circ \phi_1$, where ϕ_0 is a composition of right handed Dehn twists and ϕ_1 is some number of negative Dehn twists parallel to the boundary (we can put all the boundary parallel twists at the end since they commute with the other Dehn twists). Thus by attaching two handles to X along M in a symplectic way (see [6, 14, 30]) we can get a symplectic manifold with a boundary component (M'', ξ'') , such that (M'', ξ'') is supported by an open book with monodromy $\phi \circ \phi_0$ which is, say, k positive Dehn twists parallel to the boundary. Topologically the manifold is shown in the Figure 1. Now for each one handle in the figure add a two handle that runs over it once





FIGURE 1. On the left hand side we show an open book with k positive Dehn twists parallel to the boundary. The right hand side is homeomorphic to the left hand side.

(this can be done in a symplectic way), see Figure 2. We now have a symplectic manifold with boundary (M', ξ') . Topologically M' is obtained by $-\frac{1}{k}$ surgery on a knot in the three sphere, thus it is a homology sphere. \square

FIGURE 2. Two handle diagrams of the manifold M' .

4. TOPOLOGY OF SYMPLECTIC FILLINGS

Proof of Theorem 1.4 and 1.5. Given the hypothesized diffeomorphism ψ we can repeat the proof of Lemma 3.1 and attach two handles to X along M to form a new symplectic manifold (X', ω') with new boundary component (S^3, ξ') . Since ξ' is fillable we know it is tight. Thus ξ' is the unique tight contact structure on S^3 (see [4]). McDuff's well known extension [25] of Gromov's theorem [16] says that (X', ω') is symplectomorphic to the blow up of the standard symplectic structure on $B^4 \subset \mathbb{C}^2$. This of course implies our original symplectic filling (X, ω) only had one boundary component and that we may cap off (X', ω') by a symplectic disk bundle over S^2 to obtain a rational surface. This completes the proof of Theorem 1.5. To finish the proof of Theorem 1.4 simply note that if a rational homology sphere splits a rational surface as ours does then the component (X', ω') must be negative definite and $H_1(X; \mathbb{Z})$ must be torsional (and trivial if M is an integral homology sphere). \square

REFERENCES

- [1] F. Ding and H. Geiges, *Symplectic fillability of tight contact structures on torus bundles*, *Algebr. Geom. Topol.* **1** (2001), 153–172.
- [2] Y. Eliashberg, *A few remarks about symplectic filling*, Preprint 2003.
- [3] Y. Eliashberg, *Unique holomorphically fillable contact structure on the 3-torus*, *Internat. Math. Res. Notices* **2** (1996), 77–82.
- [4] Y. Eliashberg, *Contact 3-manifolds twenty years since J. Martinet's work*, *Ann. Inst. Fourier* **42** (1992), 165–192.
- [5] Y. Eliashberg, *On symplectic manifolds with some contact properties*, *J. Diff. Geom.* **33** (1991), 233–238.
- [6] Y. Eliashberg, *Topological characterization of Stein manifolds of dimension > 2* , *Int. J. of Math.* **1** (1990), 29–46.
- [7] Y. Eliashberg, *Filling by holomorphic discs and its applications*, *Geometry of low-dimensional manifolds*, Vol. II (Ed. Donaldson and Thomas), Cambridge, 1990.
- [8] Y. Eliashberg and W. Thurston, *Confoliations* (Amer. Math. Soc., Providence, 1998).
- [9] J. Etnyre, *Symplectic convexity in low-dimensional topology*, *Top. Appl.* **88** (1998), 3–25.
- [10] J. Etnyre and K. Honda, *On Symplectic Cobordisms*, *Math. Annalen* **323** (2002), no. 1, 31–39.
- [11] J. Etnyre and K. Honda, *Tight contact structures with no symplectic fillings*, *Invent. Math.* **148** (2002), no. 3, 609–626.
- [12] D. Gay, *Explicit concave fillings of contact three-manifolds*, *Proc. Cam. Phil. Soc.* **133** (2002), 431–441.
- [13] E. Giroux, *Géométrie de contact: de la dimension trois vers les dimensions supérieures*, *Proc. ICM-Beijing*, **2** (2002), 405–414.

- [14] R. Gompf, *Handlebody construction of Stein surfaces*, Annals of Math. **148** (1998), 619–693.
- [15] N. Goodman, *Contact Structures and Open Books*, PhD. Thesis, University of Texas Austin 2003.
- [16] M. Gromov, *Pseudo-holomorphic curves in symplectic manifolds*, Invent. math. **82** (1985), 307–347.
- [17] Kronheimer and T. Mrowka, *Witten’s conjecture and property P*, preprint 2003.
- [18] P. Kronheimer and T. Mrowka, *Monopoles and contact structures*, Invent. math. **130** (1997), 209–255.
- [19] P. Kronheimer, T. Mrowka, P. Ozsváth, and Z. Szabó. *Monopoles and lens space surgeries*, preprint 2003.
- [20] P. Lisca, *Symplectic fillings and positive scalar curvature*, *Geom. Topol.* **2** (1998), 103–116.
- [21] P. Lisca, *On symplectic fillings of 3-manifolds*, Proceedings of 6th Gökova Geometry-Topology Conference. Turkish J. Math. **23** (1999), 151–159.
- [22] P. Lisca and G. Matić, *Tight contact structures and Seiberg-Witten invariants*, Invent. Math. **129** (1997), no. 3, 509–525.
- [23] P. Lisca and A. Stipsicz, *An infinite family of tight, not semi-fillable contact three-manifolds*, preprint 2002.
- [24] D. McDuff, *Symplectic manifolds with contact type boundary*, Invent. Math. **103** (1991), 651–671.
- [25] D. McDuff, *The structure of rational and ruled symplectic 4-manifolds*, J. Amer. Math. Soc. **3** (1990), no. 3, 679–712.
- [26] H. Ohta and K. Ono, *Simple singularities and topology of symplectically filling 4-manifold*, Comment. Math. Helv. **74** (1999), 575–590.
- [27] P. Ozsváth and Z. Szabó, *Holomorphic disks and genus bounds*, preprint 2003
- [28] A. Stipsicz, *On the geography of Stein fillings of certain 3-manifolds*, Michigan Math. J. **51** (2003), no. 2, 327–337.
- [29] W. P. Thurston and H. Winkelnkemper, *On the existence of contact forms*, Proc. Amer. Math. Soc. **52** (1975), 345–347.
- [30] A. Weinstein, *Contact surgery and symplectic handlebodies*, Hokkiado Math. Journal **20** (1991), 241–251.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PENNSYLVANIA, 209 SOUTH 33RD ST., PHILADELPHIA, PA 19104-6395

E-mail address: etnyre@math.upenn.edu

URL: <http://math.upenn.edu/~etnyre>