

OPEN PROBLEMS IN GEOMETRY OF CURVES AND SURFACES

MOHAMMAD GHOMI



Date: Last revised September 2, 2019.

Research of the author was supported in part by NSF Grant DMS-1308777.

There are problems to whose solution I would attach an infinitely greater importance than to those of mathematics, for example touching ethics, or our relation to God, or concerning our destiny and our future; but their solution lies wholly beyond us and completely outside the province of science.

—Karl Friedrich Gauss (1777-1855), Quoted in J.R. Newman, *The World of Mathematics*

It is difficult and often impossible to judge the value of a problem correctly in advance; for the final award depends upon the gain which science obtains from the problem. Nevertheless we can ask whether there are general criteria which mark a good mathematical problem. An old French mathematician said: “A mathematical theory is not to be considered complete until you have made it so clear that you can explain it to the first man whom you meet on the street.” This clearness and ease of comprehension, here insisted on for a mathematical theory, I should still more demand for a mathematical problem if it is to be perfect; for what is clear and easily comprehended attracts, the complicated repels us.

—David Hilbert, Address to the International Congress of Mathematicians, 1900

Our Euclidean intuition, probably, inherited from ancient primates, might have grown out of the first seeds of space in the motor control systems of early animals who were brought up to sea and then to land by the Cambrian explosion half a billion years ago. Primates brain had been lingering for 30-40 million years. Suddenly, in a flash of one million years, it exploded into growth under relentless pressure of the sexual-social competition and sprouted a massive neocortex (70% neurons in humans) with an inexplicable capability for language, sequential reasoning and generation of mathematical ideas. Then Man came and laid down the space on papyrus in a string of axioms, lemmas and theorems around 300 B.C. in Alexandria.

—Misha Gromov, *Spaces and Questions*, 1999

I only get frightened—and it happens very rarely—when I think I have an idea.

—J. Robert Oppenheimer, *Interview with R. Murrow*, 1955

The life so short, the craft so long to learn.

—Chaucer, *Parlement of Foules*, 1382

About the cover: Detail from *The School of Athens* by Raphael, 1511

ABSTRACT. We collect dozens of well-known and not so well-known fundamental unsolved problems involving low dimensional submanifolds of Euclidean space. The list includes selections from differential geometry, Riemannian geometry, metric geometry, discrete or polyhedral geometry, geometric knot theory, theory of convex bodies, and integral geometry. The common thread through these selections are the simplicity and intuitive nature of the questions. Extensive bibliography and historical background are included for each set of problems.

CONTENTS

0. Introduction	3
1. Isometric Embeddings	4
2. Spherical Images of Surfaces	8
3. Unfoldability of Convex Polyhedra	10
4. Area and Volume of Convex Surfaces	12
5. Extremal Problems for Space Curves	12
6. Minimal and CMC Surfaces	14
7. Negatively Curved Surfaces	14
8. Umbilic Points	15
References	16

0. INTRODUCTION

Here we discuss a number of unsolved problems in geometry of curves and surfaces which have been of interest to the author over the years. Needless to say, this selection reflects the author's personal taste and (limited) perspective, although an effort has been made to include some of the oldest and best known problems in the field. Virtually all of these questions are concerned with objects in the Euclidean plane or 3-dimensional space, although in some cases higher dimensional analogues are discussed as well, while in other cases the problems are essentially intrinsic or independent of the ambient space.

0.1. Why study curves and surfaces? Curves and surfaces are to geometry what numbers are to algebra. They form the basic ingredients of our visual perception and inspire the development of far reaching mathematical tools. Yet despite centuries of pure study, not to mention a wealth of growing applications in science and technology, there are still numerous open problems in this area which are strikingly intuitive and elementary to state, pointing to fundamental gaps in our conceptions of space and shape. According to Ezra Pound, "music begins to atrophy when it departs too far from the dance" and "poetry begins to atrophy when it gets too far from music". Perhaps the same can be said of geometry, and indeed all of mathematics, if it loses sight of its natural building blocks and connections to the physical world.

0.2. Other sources for problems. There are many lists of problems in geometry and closely related fields. For a wide range of questions in differential, Riemannian and metric geometry see Yau [166, 190], Donaldson [48], and Gromov [82, 83, 84]. Some classical problems in differential geometry and many references may be found in books by Berger [21], and Burago and Zalgaller [191]. For problems involving geodesics see Burns and Matveev [28]. A large collection of problems in discrete and convex geometry are contained in the book of Croft, Falconer, and Guy [42]; also see Gardner [58] for problems involving convex bodies. For problems in minimal surface theory see the lists by Meeks [125, 126, 129], and for surfaces of constant mean curvature see Lopez [117]. A vast collection of problems in low dimensional topology is maintained by Kirby [107]. For some problems in geometric knot theory see Adams [4].

0.3. Useful websites. There are a number of websites for tracking geometric problems. The [Open Problems Project](#) [45], maintained by Demaine, Mitchell, O'Rourke, contains a wealth of problems in discrete and computational geometry. There are also growing lists of geometric problems on [Wikipedia's Unsolved Problems](#) [1] page. Some open problem in low dimensional topology are maintained at the [Low Dimensional Topology](#) [3] page. Finally numerous problems in all aspects of geometry are continually discussed on [MathOverflow](#) [2].

0.4. The earlier draft of this article. This article is a major expansion and revision of an earlier list of problems [64] which the author had collected in the Fall of 2004 for his students in a class on differential geometry in the MASS program at Penn State University. Since then several of the problems in that list have been solved: John Pardon [146] solved Gromov's question on the distortions of knots [64, Prob. 5]; Wegner [186] found examples of bicycle curves with rotation number, or relative density, different from $1/2$ [64, Prob. 1], Wilmore's conjecture [64, Prob. 18] was solved by Coda and Neves [122], and the converse to the four vertex theorem for simple closed curves [64, Prob. 18] was proved by Dahlberg [43, 47].

1. ISOMETRIC EMBEDDINGS

A fundamental theme in surface theory, which dates back to the development of this subject by Gauss [59], is determining the extent to which the intrinsic geometry or metric structure of a surface contributes to its global shape in the ambient space. Questions of this type, which we discuss below may be phrased within the context of the uniqueness and existence of isometric embeddings of 2-dimensional Riemannian manifolds in Euclidean 3-space \mathbf{R}^3 .

1.1. Flexibility of closed surfaces. As every child soon learns, an egg shell is not flexible. The mathematical reason is the rigidity theorem for convex surfaces first proved by Cauchy for polyhedra, then extended to smooth surfaces by Hilbert, Cohn-Vossen, Weyl, Nirenberg, and finally generalized to all convex surfaces by Alexandrov and Pogorelov [94, 9]. This theorem states that isometric (closed) convex surfaces are congruent, or, to put it more succinctly, convex surfaces are (isometrically) *rigid*. Finding an analogue of this result for general (nonconvex) surfaces

is one of the oldest problems in geometry [190], [188, Problem 50], which may be traced back to Euler [54, p. 494–496] for polyhedral surfaces, see [76, 78, 115], and Maxwell [124] for smooth surfaces; however, to quote Chern [32, p. 211], “practically nothing is known” about it.

To state the problem explicitly, let us say that a surface is (isometrically) *flexible*, within a given smoothness class, if it admits a continuous deformation which preserves its smoothness and does not change its intrinsic metric. For instance, a sheet of paper is flexible in the class of C^∞ surfaces, since it can be rolled into a cylinder. It is also well-known, though not all obvious, that a spherical cap (a piece of sphere cut off by a plane) is smoothly flexible. No one, however, has ever found a flexible *closed* surface in the smoothness class C^2 , or even $C^{1,1}$, where by closed we mean compact without boundary. Hence the following question:

Problem 1.1 (Euler [54], 1776; and Maxwell [124], 1819). *Does there exist a closed C^2 surface in Euclidean space \mathbf{R}^3 which is flexible?*

Note that flexibility is a stronger notion than nonrigidity. Indeed it is easy to construct C^∞ surfaces which are not rigid: consider for instance a closed surface with a dimple which is surrounded by a flat rim, and replace the dimple by its reflection. Such a transformation, however, cannot be carried out continuously for it would force the mean curvature vector to switch sides. One can even construct closed *analytic* surfaces which are not rigid [151], though these constructions are far more subtle.

Secondly, it is important to note that the answer to the analogue of Problem 1.1 in the category of polyhedral surfaces is yes! The first example of a flexible closed polyhedral surface was constructed by Robert Connelly [39]; however, these flexible polyhedra are not so natural. Indeed Gluck [76] had shown that almost all closed polyhedral surfaces are rigid, see the book of Igor Pak [145]. Further, Stoker [178] explicitly described a large class of nonconvex polyhedral surfaces which are rigid, which were further studied and generalized by Rodriguez and Rosenberg [153]. Stoker polyhedra have vertices which are either convex or are of saddle type. The latter means that the vertex has degree 4 and the edges around that vertex alternate between valley and ridge types. Saddle vertices have the property that unit normals to the faces around the vertex form the vertices of a convex geodesic polygon in the sphere. This local convexity property of the Gauss map is the key ingredient to generalizing Cauchy’s rigidity theorem to nonconvex polyhedra [153]. Rigidity of some other classes of nonconvex polyhedra have also been established in papers of Schlenker with Izmetsiev [103] and Connolly [40].

An even greater surprise, than flexibility of nonconvex polyhedra, is the flexibility of C^1 isometric embeddings. For instance one can squeeze the unit sphere into an arbitrarily small ball through a continuous one-parameter family of isometric C^1 immersions [23]! Thus the C^2 requirement in the above problem is not superfluous. The remarkable nature of C^1 isometric embeddings were first discovered by Nash and Kuiper, and further developed by Gromov [81] within the frame work of the *h*-Principle theory; see also the book by Eliashberg and Mishachev [53]. In this

context, the flexibility of the C^1 isometric embeddings corresponds to the existence of a parametric h -principle for these maps.

The fourth point to keep in mind, with regard to Problem 1.1, is that it is important that the ambient space be \mathbf{R}^3 . Indeed Pinkall [147] showed that flat tori flex isometrically in \mathbf{S}^3 —a phenomenon which was studied further by Kitagawa [108].

Finally we should mention the paper by Almgren and Rivin [12] where it is shown that the mean curvature integral must be preserved under isometric flexing.

1.2. Rigidity of tight surfaces. Traditional approaches for studying Problem 1.1 consist of breaking the surface up into negatively curved and positively curved regions, and studying the flexibility of these regions individually. This method was used by Alexandrov [7] to establish the rigidity of analytic surfaces with total positive curvature 2π (e.g., tori of revolution) which are also known as *tight surfaces* [33, 113, 29], and are natural higher genus analogues of convex surfaces. Subsequently, Nirenberg [140, 94] made a beautiful but incomplete attempt at generalizing Alexandrov’s result to smooth surfaces. Thus we have

Problem 1.2 (Alexandrov [7] 1938; and Nirenberg [140] 1962). *Are all smooth tight surfaces in \mathbf{R}^3 rigid?*

In the polyhedral case Banchoff [17] has shown that the answer to the above question is no. Nirenberg observed that the convex part of smooth tight surfaces are rigid, thus the above problem comes down to the following question:

Problem 1.3. *Are negatively curved annuli bounded by a pair of fixed convex planar curves rigid?*

The main obstacle Nirenberg encountered in answering this question could be lifted by a negative answer to the following question: *Can a negatively curved annulus in \mathbf{R}^3 , bounded by a pair of convex planar curves, have a closed asymptotic curve?* A curve in a surface is *asymptotic* if the normal curvature of the surface always vanishes in directions tangent to the curve. One way to approach the above problem is as follows. Suppose that we have a smooth closed curve Γ immersed in \mathbf{R}^3 which admits a continuous family of osculating planes, i.e., planes which contain the first and second derivatives of Γ . Then the binormal vector field of Γ may be defined as any continuous vector field which is orthogonal to these osculating planes. If Γ is an asymptotic curve in a surface M , then the gauss map of M gives a binormal vector field along Γ . Thus Nirenberg’s problem leads to the following question:

Problem 1.4. *Let Γ be a smooth closed curve immersed in \mathbf{R}^3 . Suppose that Γ has a continuous binormal vector field B which is one-to-one. Does it follow then that the ribbon (Γ, B) is twisted? In other words, must the linking number between Γ and $\Gamma + \epsilon B$ be nonzero?*

A positive answer to the last question will settle Problem 1.2. This approach has been discussed in [112]; however, subject to regularity assumptions not warranted by the question. Further, existence of closed asymptotic curves on negatively curved

surfaces has also been studied by Arnold [16] who raised a number of interesting questions as well.

1.3. Surfaces with prescribed boundary. On the other hand, understanding the flexibility of positively curved surfaces with boundary is also important as far as Problems 1.1 and 1.2 are concerned. Here, a fundamental questions is:

Problem 1.5 (Yau [189], 1990). *Given a metric of positive curvature on the disk what is the condition on a space curve to form the boundary of an isometric embedding of the disk?*

A nontrivial *necessary* condition has been discovered by H. Rosenberg [157], involving the self-linking number, and a result of the author [61] provides a *sufficient* criterion; however, a complete characterization is not yet known. In particular, Gluck and Pan [77] constructed an example to show that Rosenberg's condition was not sufficient. Also we should note that Guan and Spruck [87], as well as Trudinger and Wang [184], have shown that if a curve bounds a surface of positive curvature, then it bounds a surface of *constant* positive curvature. Thus Problem 1.5 may be regarded as a boundary value problem for a PDE of Monge-Ampere type. Another question related to Problem 1.5 is:

Problem 1.6 (Rosenberg [157], 1993). *Does every curve bounding a surface of positive curvature in 3-space have (at least) four points where the torsion vanishes?*

The author has shown recently that the answer to the last question is yes when the surface is topologically a disk [72], through a comprehensive study of the structure of convex caps in locally convex surfaces. Convex caps play a major role in the seminal works of Alexandrov [10] and Pogorelov [149] on the isometric embeddings of convex surfaces, as well as in other fundamental results in this area such as the works by van Heijenoort [185], Sacksteder [160], and Volkov [9, Sec. 12.1]. In these studies, however, the underlying Riemannian manifolds are assumed to be complete, or nearly complete, as in the works of Greene and Wu [79, 80]. In [72], on the other hand, the author studies caps of manifolds with boundary, as in the author's previous work with Alexander [5] and Alexander and Wong [6].

The author's solution to the last problem in the genus zero case establishes a far reaching generalization of the classical four vertex theorem for planar curves. The study of special points of curvature and torsion of closed curves has generated a vast and multifaceted literature since the works of Mukhopadhyaya [135] and A. Kneser [111] on vertices of planar curves were published in 1910–1912, although aspects of these investigations may be traced even further back to the study of inflections by Möbius [133] and Klein [109], see [69, 68]. The first version of the four vertex theorem for space curves, which was concerned with curves lying on smooth strictly convex surfaces, was stated by Mohrmann [134] in 1917, and proved by Barner and Flohr [19] in 1958. This result was finally extended to curves lying on the boundary of any convex body by Sedykh [169] in 1994, after partial results by other authors [141, 22], see also Romero-Fuster and Sedykh [154] for further refinements.

Among various applications of four vertex theorems, we mention a paper of Berger and Calabi et al. [150] on physics of floating bodies, and recent work of Bray and Jauregui [26] in general relativity. See also the works of Arnold [14, 15] for relations with contact geometry, the book of Ovsienko and Tabachnikov [144] for projective geometric aspects, Angenent [13] for connections with mean curvature flow, which are also discussed in [68], and Ivanisvili et al. [100, 101] for applications to the study of Bellman functions. Other references and more background on four vertex theorems may be found in [47, 181, 67, 145, 75].

1.4. Rigidity of Punctured surfaces. Greene and Wu showed that smooth convex surfaces with finitely many points deleted remain rigid [79, 80]. So it is natural to wonder:

Problem 1.7. *Are there some nonconvex surfaces which remain rigid after finitely many points of them have been deleted. For instance, are punctured analytic tight surfaces, such as a torus of revolution, rigid?*

1.5. Existence problems. By a celebrated theorem of Nash every Riemannian manifold admits an isometric embedding into a Euclidean space of sufficiently high dimension; however, much remains unknown about the smallest dimension where this is possible for a given class of manifolds [81, 84], including 2-dimensional manifolds or surfaces.

Problem 1.8 (The global isometric embedding problem, Yau [189] 1993; Gromov [82]). *Can every C^∞ 2-dimensional Riemannian manifold be isometrically embedded in \mathbf{R}^4 ?*

It is well-known that locally the answer to the above question is yes; however, the main local problem is the following:

Problem 1.9 (The local isometric embedding problem). *Given a C^∞ metric in a neighborhood of a point in a 2-dimensional Riemannian manifold, does there exist an isometric embedding of some neighborhood of that point into \mathbf{R}^3 ?*

The last problem has been well studied, see the book by Han and Hong for references [94]. In particular, it is well-known that the answer to the last problem is yes if the curvature is strictly positive, or strictly negative. Further, in the cases where the curvature “changes sign cleanly”, i.e., 0 is a regular value of the curvature function, it is also known that the answer is yes. On the other hand, Pogorelov [148] constructed a C^2 metric which does not admit a local isometric embedding in \mathbf{R}^3 . See also the paper by Nadirashvili and Yuan [137] for more results in this direction.

2. SPHERICAL IMAGES OF SURFACES

Most problems in surface theory, including the rigidity problems mentioned earlier, may be rephrased or studied in terms of the unit normal vector field or the Gauss map of surfaces. Here we discuss three problems in this area, which may also be classified as questions in convex integration theory.

2.1. Directed immersions. To every (\mathcal{C}^1) immersion $f: M^n \rightarrow \mathbf{R}^{n+1}$ of a closed oriented n -manifold M , there corresponds a unit normal vector field or *Gauss map* $G_f: M \rightarrow \mathbf{S}^n$, which generates a set $G_f(M) \subset \mathbf{S}^n$ known as the *spherical image* of f . Conversely, one may ask:

Problem 2.1 (Gromov [82]). *For which sets $A \subset \mathbf{S}^n$ is there an immersion $f: M \rightarrow \mathbf{R}^{n+1}$ such that $G_f(M) \subset A$?*

Such a mapping would be called an *A-directed immersion* of M [53, 81, 158, 177]. It is well-known that when $A \neq \mathbf{S}^n$, f must have double points, and M must be parallelizable, e.g., M can only be the torus \mathbf{T}^2 when $n = 2$. Furthermore, the only known necessary condition on A is the elementary observation that $A \cup -A = \mathbf{S}^2$, while there is also a sufficient condition due to Gromov [81, Thm. (D'), p. 186]: *$A \subset \mathbf{S}^n$ is open, and there is a point $p \in \mathbf{S}^n$ such that the intersection of A with each great circle passing through p includes a (closed) semicircle.* Note that, when $n \geq 2$, examples of sets $A \subset \mathbf{S}^n$ satisfying this condition include those which are the complement of a finite set of points without antipodal pairs. Thus the spherical image of a closed hypersurface can be remarkably flexible. Like most h -principle or convex integration type arguments, however, the proof does not yield specific examples. It is therefore natural to ask [81]: “Is there a ‘simple’ immersion $\mathbf{T}^2 \rightarrow \mathbf{R}^3$ whose spherical image misses the four vertices of a regular tetrahedron in \mathbf{S}^2 ?” The author has given an affirmative answer to this question [66], and more generally has given a short constructive proof of the sufficiency of a slightly stronger version of Gromov’s condition mentioned above for the existence of A -directed immersions of parallelizable manifolds $M^{n-1} \times \mathbf{S}^1$, where M^{n-1} is closed and orientable.

2.2. The shadow problem. Next we describe a problem which involves the notion of *shadows* or *shades* on illuminated surfaces and has applications to the so called “shape from shading” problems in computer vision [98], and various problems in calculus variations [34, 60]. Let $f: M \rightarrow \mathbf{R}^{n+1}$ be a closed oriented n -dimensional hypersurface in Euclidean space as before and $G_f: M \rightarrow \mathbf{S}^n$ be its Gauss map. Then for every unit vector $u \in \mathbf{S}^n$ (corresponding to the direction of light) the *shadow* (or *shade* as it is known in computer vision [98]) is defined as

$$(1) \quad S_u := \{p \in M \mid \langle G_f(p), u \rangle > 0\},$$

where $\langle \cdot, \cdot \rangle$ is the standard innerproduct. In 1978, motivated by problems concerning the stability of constant mean curvature (CMC) surfaces (or soap films) [187], H. Wente formulated the following question, which has since become known as the “shadow problem:

Problem 2.2 (Wente [187], 1978). *Does connectedness of the shadows imply that $f(M)$ is convex?*

In [63] the author found a complete solution to this problem in 3-space: In \mathbf{R}^3 , the answer to Problem 2.2 is yes provided that each shadow is *simply* connected, or that M is topologically a sphere; otherwise, the answer is no. In particular, there is a smooth embedded topological torus in \mathbf{R}^3 which has connected shadows in every

direction. Indeed as the author later showed [65], every closed orientable 2-manifold admits a smooth embedding in \mathbf{R}^3 which has connected shadows in every direction, thus disproving a conjecture of J Choe [34]. But there is nothing known in higher dimensions.

2.3. Minkowski problem for nonconvex surfaces. The famous problem of Minkowski, which has been completely solved [165], is concerned with proving that every convex surface is uniquely determined by its curvature prescribed on the sphere via the Gauss map. The discrete version of this problem states that two convex polytopes whose faces are parallel and have the same area are congruent. As has been pointed out by Yau [188] it would be interesting to find analogues of these results for nonconvex surfaces:

Problem 2.3. *Let $M, M' \subset \mathbf{R}^3$ be smooth orientable closed surfaces. Suppose there exists a diffeomorphism $f: M \rightarrow M'$ which preserved the Gauss curvature and the Gauss map. Does it follow then that M and M' are congruent?*

This problem also has a discrete version:

Problem 2.4. *Let $P, P' \subset \mathbf{R}^3$ be polyhedral surfaces. Suppose that the faces of P and P' are parallel and have the same area. Does it follow then that P and P' are congruent?*

3. UNFOLDABILITY OF CONVEX POLYHEDRA

A well-known problem [46, 142, 145, 195], which may be traced back to the Renaissance artist Albrecht Dürer [49], is concerned with cutting a convex polyhedral surface P along some spanning tree T of its edges so that it may be isometrically embedded, or unfolded without overlaps, into the plane. If P admits such a one-to-one unfolding (for some choice of T) then we say that P is *unfoldable*.

Problem 3.1 (Dürer [49], 1525; Shephard [170], 1975). *Is every convex polytope unfoldable?*

The author has shown [70] that this is always possible after an affine transformation of the surface. In particular, unfoldability of a convex polyhedron does not depend on its combinatorial structure, which settles a problem of Croft, Falconer, and Guy [41, B21]. To describe this result more explicitly, assume that the polyhedron P is in *general position* with respect to L , by which we mean that for a unit vector u parallel to L the height function $\langle \cdot, u \rangle$ has a unique maximizer and a unique minimizer among the vertices of P . There is an open dense set of such lines in the real projective space \mathbf{RP}^2 . In [70] it is shown that P becomes unfoldable after a sufficiently large stretching in any such direction. To describe what we mean by *stretching* it would be convenient to assume, after a rotation, that L is parallel to the z -axis in \mathbf{R}^3 , then our transformation is given by $(x, y, z) \mapsto (x, y, \lambda z)$. This will have the effect of making P arbitrarily “thin” or “needle-shaped”. Thus, roughly speaking, needle-shaped convex polytopes are unfoldable.

Dürer’s problem is the most famous question in the theory of foldings and unfoldings [46], or origami research, which has had wide applications ranging from

assembling satellite dishes in space to the expansion of implanted stents in human arteries. For extensive background on this problem, and an outline of the author's solution in the affine case, see

people.math.gatech.edu/~ghomi/Talks/durerslides.pdf

Yet there is still no algorithm for unfolding a convex polytope injectively despite intense efforts by computer scientists. The difficulty of Dürer's problem is that it is essentially an intrinsic question, yet there is no known intrinsic method to detect the edges of a convex polyhedron. Indeed, Alexandrov's embedding theorem for convex surfaces [9]—which states that any locally convex polyhedral metric on \mathbf{S}^2 may be realized as a convex polyhedron in \mathbf{R}^3 —is not constructive and gives no hint as to which geodesics between a pair of vertices are realized as edges; see also Pak [145]. In 2008, a more constructive proof was given by Bobenko and Izestiev [24]; however, this proof does not specify the location of the edges either. Thus Dürer's problem provides another opportunity to deepen our understanding of isometric embeddings.

Problem 3.2. *Does there exist a reasonably simple algorithm for detecting the edges of a convex polyhedron intrinsically?*

The edge graph of P is not the unique graph in P whose vertices coincide with those of P , whose edges are geodesics, and whose faces are convex. It seems reasonable to expect that Dürer's conjecture should be true if and only if it holds for this wider class of *pseudo edge graphs*. This approach was studied by Tarasov [180] in 2008, who announced some negative results in this direction; however, Tarasov's paper has not been published. Checking the validity of Tarasov's constructions would be an important step towards solving Dürer's problem:

Problem 3.3. *Does there exist a convex polyhedron with a pseudo edge graph which is not unfoldable.*

The earliest known examples of simple edge unfoldings for convex polyhedra are due to Dürer [49], although the problem which bears his name was first formulated by Shephard [170]. Furthermore, the assertion that a solution can always be found, which has been dubbed Dürer's conjecture, appears to have been first published by Grünbaum [85, 86]. There is empirical evidence both for and against this supposition. On the one hand, computers have found simple edge unfoldings for countless convex polyhedra through an exhaustive search of their spanning edge trees. On the other hand, there is still no algorithm for finding the right tree [164, 118], and computer experiments suggest that the probability that a random edge unfolding of a generic polyhedron overlaps itself approaches 1 as the number of vertices grow [163]. General cut trees have been studied at least as far back as Alexandrov [9] who first established the existence of simple unfoldings (not necessarily simple *edge* unfoldings) for all convex polyhedra, see also [99, 131, 44] for recent related results. Other references and background may be found in [46].

4. AREA AND VOLUME OF CONVEX SURFACES

Problem 4.1 (A. D. Alexandrov [8]). *Of all convex surfaces with a fixed intrinsic diameter, is the one with the greatest area a doubled disk?*

It is known that the answer to the above question is affirmative for surfaces of revolution [121] and that in the class of tetrahedra the maximizer is, rather surprisingly, the regular tetrahedron [119]. See also [120] for results relating the intrinsic and extrinsic diameter of convex surfaces. For other partial results using techniques in Riemannian geometry see [172, 161]

Problem 4.2 (Volume of surfaces of constant width). *Let $S \subset \mathbf{R}^3$ be a closed surface of constant width and fixed area. How small can the volume of S be?*

The above problem has been solved in \mathbf{R}^2 : the Reuleaux triangle has the least area among all closed curves of constant width. For a recent proof of this result, which is originally due to Blaschke and Lebesgue, see Harrell [95]. For other references see [42].

Problem 4.3 (Surfaces with strips of constant area). *Let $S \subset \mathbf{R}^3$ be a closed surface of diameter d . Suppose that there exists a constant $h < d$ so that whenever a pair of planes separated by a distance of h intersect S , the area of S contained between these planes is constant. Does it then follow that S is a sphere?*

5. EXTREMAL PROBLEMS FOR SPACE CURVES

What is the smallest length of wire which can be bent into a shape that never falls through the gap behind a desk? What is the shortest orbit which allows a satellite to survey a spherical asteroid? These are well-known open problems [193, 143, 42, 96] in classical geometry of space curves $\gamma: [a, b] \rightarrow \mathbf{R}^3$, which are concerned with minimizing the length L of γ subject to constraints on its *width* w and *inradius* r respectively. Here w is the infimum of the distances between all pairs of parallel planes which bound γ , while r is the supremum of the radii of all spheres which are contained in the convex hull of γ and are disjoint from γ . In 1994–1996 Zalgaller [192, 193] conjectured four explicit solutions to these problems, including the cases where γ is restricted to be closed, i.e., $\gamma(a) = \gamma(b)$. Recently the author has been able to employ a combination of integral geometric and topological techniques to confirm Zalgaller’s conjectures [71] between 83% and 99% of their stated value, while he has also found a counterexample to one of them, but sharp answers have not yet been found.

Problem 5.1 (Zalgaller [192] 1994). *What is the shortest curve in \mathbf{R}^3 with a given width or inradius?*

The author has shown that, for any rectifiable curve $\gamma: [a, b] \rightarrow \mathbf{R}^3$,

$$(2) \quad \frac{L}{w} \geq 3.7669.$$

Furthermore if γ is closed,

$$(3) \quad \frac{L}{w} \geq \sqrt{\pi^2 + 16} > 5.0862.$$

In [192] Zalgaller constructs a curve, “ L_3 ”, with $L/w \leq 3.9215$. Thus (2) is better than 96% sharp (since $3.7669/3.9215 \geq 0.9605$). Further, there exists a closed cylindrical curve with $L/w < 5.1151$, which shows that (3) is at least 99.43% sharp. In particular, the length of the shortest closed curve of width 1 is approximately 5.1.

The author has also found estimates for the inradius problem. Obviously $w \geq 2r$, and thus the above inequalities immediately yield $L/r \geq 7.5338$ for general curves, and $L/r \geq 10.1724$ for closed curves. Using different techniques, however, these estimates may be improved as follows: for any rectifiable curve $\gamma: [a, b] \rightarrow \mathbf{R}^3$,

$$(4) \quad \frac{L}{r} \geq \sqrt{(\pi + 2)^2 + 36} > 7.9104.$$

Furthermore if γ is closed,

$$(5) \quad \frac{L}{r} \geq 6\sqrt{3} > 10.3923.$$

In [194, Sec. 2.12] Zalgaller constructs a spiral curve with $L/r \leq 9.5767$, which shows that (4) is better than 82.6% optimal. Further, in [193], he produces a curve composed of four semicircles with $L/r = 4\pi$; see also [143] where this “baseball stitches” curve is rediscovered in 2011. Thus we may say that (5) is better than 82.69% optimal.

Both the width and inradius problems may be traced back to a 1956 question of Bellman [20] motivated by harmonic analysis: how long is the shortest escape path for a random point (lost hiker) inside an infinite parallel strip (forest) of known width? See [56] for more on these types of problems. Our width problem is the analogue of Bellman’s question in \mathbf{R}^3 . The inradius problem also has an intuitive reformulation known as the “sphere inspection” [194, 143] or the “asteroid surveying” problem [31]; see

mathoverflow.net/questions/69099/shortest-closed-curve-to-inspect-a-sphere

To describe this variation, let us say that a space curve γ *inspects* the sphere \mathbf{S}^2 , or is an *inspection curve*, provided that γ lies outside \mathbf{S}^2 and for each point x of \mathbf{S}^2 there exists a point y of γ such that the line segment xy does not enter \mathbf{S}^2 (in other words, x is “visible” from y). It is easy to see that γ inspects \mathbf{S}^2 , after a translation, if and only if its inradius is 1 [193, p. 369]. Thus finding the shortest inspection curve is equivalent to the inradius problem for $r = 1$.

Problem 5.2 (Volume of the convex hull of closed curves). *Let Γ be a closed curve of fixed length L in \mathbf{R}^3 . How big can the volume of the convex hull of Γ be.*

For a partial result under symmetry conditions for the above problem see [130]. The above problem has been solved in Euclidean spaces of even dimensions [168]. Also the problem is solved for open arcs in \mathbf{R}^3 [51]. For some other related results and questions see [193].

Problem 5.3 (Area of the convex hull of closed curves). *Let Γ be a closed curve of fixed length L in \mathbf{R}^3 , and A be the area of the convex hull of Γ . Show that A is biggest when Γ is a circle, in which case we consider the convex hull of Γ as a doubly covered disk.*

If Γ is a simple curve which lies on the boundary of its convex hull, then it divides the boundary of the convex hull into a pair of disks each of which have zero curvature. This observation together with the isoperimetric inequality for surfaces of nonpositive curvature, first proved by Andre Weil, may be used to solve the above problem in the case where Γ is simple and lies on the boundary of its convex hull.

6. MINIMAL AND CMC SURFACES

Problem 6.1 (Meeks [125, 129, 126], 1978). *Is every compact connected minimal surface bounded by a pair of convex planar curves topologically an annulus?*

An affirmative solution to the above problem would lead to a generalization of Shiffman's classical theorem [171] on level set of minimal annuli. Partial or related results have been obtained in [167, 127, 128, 138, 155, 104, 52, 62]. See the article by Hoffman and Meeks [97] for a nice introduction to the conjecture.

Problem 6.2 (Earp, Fabiano, Meeks, and Rosenberg [50], 1991). *Does there exist an embedded compact surface of constant mean curvature which is bounded by a circle, but is not a piece of a sphere.*

If the surface is only immersed, not embedded, then the answer to the last problem is yes, as was shown by Kapouleas [106]. Furthermore [11] Alias, Lopez, and Palmer showed that the answer is yes, if the surface is assumed to be stable and a topological disk.

Problem 6.3 (Ros and Rosenberg [156], 1996). *Show that any compact embedded CMC surface which is bounded by a convex planar curve, and lies on one side of the boundary plane, is topologically a disk.*

The above problem has been affirmatively solved by Barbosa and Jorge assuming that the surface is stable [18]. For more background and references related to the last problem see the book of Lopez [117].

7. NEGATIVELY CURVED SURFACES

The next problem is due to John Milnor [132], and would generalize famous theorems of Hilbert and Efimov. Hilbert showed that there exists no complete surfaces of constant negative curvature in \mathbf{R}^3 , and Efimov proved that there exists no complete surfaces of negative curvature in \mathbf{R}^3 whose curvature is bounded away from zero. Proof of Hilbert's theorem may be found in many elementary texts on differential geometry. For Efimov's proof see [132]. Some relatively recent proofs of these results have been announced in [152].

Problem 7.1 (Milnor [132], 1972). *Are there any complete surfaces of negative curvature in Euclidean 3-space whose principal curvatures are bounded away from zero?*

An even older problem on negatively curved surfaces is the following:

Problem 7.2 (Hadamard [90, 27, 159], 1898). *Are there any complete negatively curved surfaces embedded in the unit ball?*

In the immersed case, this conjecture was settled by Nadirashvili [136] who constructed complete minimal negatively curved surfaces in a ball, see also [105, 36, 123, 116]; however, such surfaces cannot be embedded, since by a result of Colding and Minicozzi [35], an embedded minimal surface must be unbounded. An approach for constructing a possible counterexample to the above conjecture has been outlined by Rozendorn [159].

Problem 7.3. *Does there exist any complete negatively curved surfaces with negative Euler characteristic contained in between a pair of parallel planes in \mathbf{R}^3 .*

It is not difficult to construct such surfaces with nonnegative Euler characteristic, see the author's paper with Chris Connell [37, Note 1.4]. See also [38] and [30] for relevant results on topology of negatively surfaces.

8. UMBILIC POINTS

Problem 8.1 (Carathéodory, 1922). *Show that every closed convex surface in \mathbf{R}^3 has at least two umbilic points.*

According to Struik [179], the earliest references to the conjecture attributed to Carathéodory appear in the works of Cohn-Vossen, Blaschke, and Hamburger dating back to 1922. The first significant results on the conjecture were due to Hamburger, who established the analytic case in a series of long papers [91, 92, 93] published in 1940–41. Attempts to find shorter proofs attracted the attention of Bol [25], Klotz [110], and Titus [182] in the ensuing decades. As late as 1993, however, Scherbel [162] was still correcting some errors in these simplifications, while reconfirming the validity of Hamburger's theorem. Another reexamination of the proof of the analytic case appears in a comprehensive paper of Ivanov [102] who supplies his own arguments for clarifying various details. All the works mentioned thus far have been primarily concerned with establishing the analytic version of a local conjecture attributed to Loewner, which states that

Problem 8.2 (Loewner). *Show that the index of any singularity of a principal line fields on a surface is at most one.*

A positive resolution of Loewner's conjecture would imply Carathéodory's conjecture via Poincaré-Hopf index theorem. See Smyth and Xavier [173, 174, 175] for studies of Loewner's conjecture in the smooth case, and Lazarovici [114] for a global result on principal foliations. Another global result is by Feldman [55], who showed that generic closed convex surfaces have four umbilics; also see [74] for some

applications of the h -principle to studying homotopy classes of principal lines. A global generalization of Carathéodory's conjecture is discussed in [57]. There is also an interesting analogue of the conjecture for noncompact complete convex hyper-surfaces:

Problem 8.3 (Toponogov [183], 1995). *Let M be a complete noncompact convex surface in \mathbf{R}^3 , with principal curvatures k_1, k_2 , then show that $\inf_M |k_1 - k_2| = 0$.*

In other words, Toponogov suggests that every complete noncompact convex surface must have an umbilic “at infinity”. A number of approaches to proving the Carathéodory or Loewner conjecture in the smooth case are discussed in [144, 139, 88], and more references or background may be found in [176, 89]. In recent joint work with Ralph Howard [73], the author has found some real evidence in support of this conjecture. At the same time we constructed a number of surfaces which may be considered counterexamples in a weak sense. A number of approaches to proving the Carathéodory or Loewner conjecture in the smooth case are discussed in [144, 139, 88]. In particular Guilfoyle and Klingenberg [88] announced a solution to the conjecture in 2008, although that work does not seem to have been published yet. More references or background may be found in [176, 89].

REFERENCES

- [1] http://en.wikipedia.org/wiki/list_of_unsolved_problems_in_mathematics.
- [2] <http://mathoverflow.net>.
- [3] <https://ldtopology.wordpress.com/open-problems/>.
- [4] C. C. Adams. *The knot book*. American Mathematical Society, Providence, RI, 2004. An elementary introduction to the mathematical theory of knots, Revised reprint of the 1994 original.
- [5] S. Alexander and M. Ghomi. The convex hull property and topology of hypersurfaces with nonnegative curvature. *Adv. Math.*, 180(1):324–354, 2003.
- [6] S. Alexander, M. Ghomi, and J. Wong. Topology of Riemannian submanifolds with prescribed boundary. *Duke Math. J.*, 152(3):533–565, 2010.
- [7] A. D. Alexandrov. On a class of closed surfaces. *Mat. Sbornik*, 4:69–77, 1938.
- [8] A. D. Alexandrov. *Die innere Geometrie der konvexen Flächen*. Akademie-Verlag, Berlin, 1955.
- [9] A. D. Alexandrov. *Convex polyhedra*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2005. Translated from the 1950 Russian edition by N. S. Dairbekov, S. S. Kutateladze and A. B. Sossinsky, With comments and bibliography by V. A. Zalgaller and appendices by L. A. Shor and Yu. A. Volkov.
- [10] A. D. Alexandrov. *A. D. Alexandrov selected works. Part II*. Chapman & Hall/CRC, Boca Raton, FL, 2006. Intrinsic geometry of convex surfaces, Edited by S. S. Kutateladze, Translated from the Russian by S. Vakhrameyev.
- [11] L. Alías, R. López, and B. Palmer. Stable constant mean curvature surfaces with circular boundary. *Proceedings of the American Mathematical Society*, 127(4):1195–1200, 1999.
- [12] F. J. Almgren, Jr. and I. Rivin. The mean curvature integral is invariant under bending. In *The Epstein birthday schrift*, volume 1 of *Geom. Topol. Monogr.*, pages 1–21 (electronic). Geom. Topol. Publ., Coventry, 1998.
- [13] S. Angenent. Inflection points, extatic points and curve shortening. In *Hamiltonian systems with three or more degrees of freedom (S'Agaró, 1995)*, volume 533 of *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.*, pages 3–10. Kluwer Acad. Publ., Dordrecht, 1999.

- [14] V. I. Arnold. *Topological invariants of plane curves and caustics*, volume 5 of *University Lecture Series*. American Mathematical Society, Providence, RI, 1994. Dean Jacqueline B. Lewis Memorial Lectures presented at Rutgers University, New Brunswick, New Jersey.
- [15] V. I. Arnold. The geometry of spherical curves and quaternion algebra. *Uspekhi Mat. Nauk*, 50(1(301)):3–68, 1995.
- [16] V. I. Arnold. Topological problems in the theory of asymptotic curves. *Tr. Mat. Inst. Steklova*, 225(Solitary Geom. Topol. na Perekrest.):11–20, 1999.
- [17] T. F. Banchoff. Non-rigidity theorems for tight polyhedra. *Arch. Math. (Basel)*, 21:416–423, 1970.
- [18] J. L. M. Barbosa and L. R. Jorge. Uniqueness of stable h -surfaces spanning $s^1(1)$. *An. Acad. Brasil. Cienc.*, 67:1–5, 1995.
- [19] M. Barner and F. Flohr. Der vierscheitesatz und seine verallgemeinerungen. *Der Mathematikunterricht*, 44:43–73, 1958.
- [20] R. Bellman. A minimization problem. *Bulletin of the AMS*, 62:270, 1956.
- [21] M. Berger. *A panoramic view of Riemannian geometry*. Springer-Verlag, Berlin, 2003.
- [22] T. Bisztriczky. On the four-vertex theorem for space curves. *J. Geom.*, 27(2):166–174, 1986.
- [23] D. Bleeker. Isometric deformations of compact hypersurfaces. *Geom. Dedicata*, 64(2):193–227, 1997.
- [24] A. Bobenko and I. Izmistiev. Alexandrov’s theorem, weighted delaunay triangulations, and mixed volumes. In *Annales de l’institut Fourier*, volume 58, pages 447–505, 2008.
- [25] G. Bol. Über Nabelpunkte auf einer Eifläche. *Math. Z.*, 49:389–410, 1944.
- [26] H. L. Bray and J. L. Jauregui. On curves with nonnegative torsion. *arXiv:1312.5171*, 2014.
- [27] Y. D. Burago and S. Z. Shefel’. The geometry of surfaces in Euclidean spaces [MR1039818 (91d:53004)]. In *Geometry, III*, volume 48 of *Encyclopaedia Math. Sci.*, pages 1–85, 251–256. Springer, Berlin, 1992.
- [28] K. Burns and V. S. Matveev. Open problems and questions about geodesics. *arXiv preprint arXiv:1308.5417*, 2013.
- [29] T. E. Cecil and P. J. Ryan. *Tight and taut immersions of manifolds*. Pitman (Advanced Publishing Program), Boston, Mass., 1985.
- [30] H. Chan. Topological uniqueness of negatively curved surfaces. *Nagoya Mathematical Journal*, 199:137–149, 2010.
- [31] T. M. Chan, A. Golynski, A. Lopez-Ortiz, and C.-G. Quimper. The asteroid surveying problem and other puzzles. In *Proceedings of the Nineteenth Annual Symposium on Computational Geometry*, SCG ’03, pages 372–373, New York, NY, USA, 2003. ACM.
- [32] S. S. Chern. The geometry of G -structures. *Bull. Amer. Math. Soc.*, 72:167–219, 1966.
- [33] S.-s. Chern and R. K. Lashof. On the total curvature of immersed manifolds. *Amer. J. Math.*, 79:306–318, 1957.
- [34] J. Choe. Index, vision number and stability of complete minimal surfaces. *Arch. Rational Mech. Anal.*, 109(3):195–212, 1990.
- [35] T. H. Colding and W. P. Minicozzi. The calabi-yau conjectures for embedded surfaces. *Annals of mathematics*, 167(1):211–243, 2008.
- [36] P. Collin and H. Rosenberg. Notes sur la démonstration de n. nadirashvili des conjectures de hadamard et calabi-yau. *Bulletin des sciences mathématiques*, 123(7):563–575, 1999.
- [37] C. Connell and M. Ghomi. Topology of negatively curved real affine algebraic surfaces. *J. Reine Angew. Math.*, 624:1–26, 2008.
- [38] C. Connell and J. Ullman. Ends of negatively curved surfaces in euclidean space. *manuscripta mathematica*, 131(3-4):275–303, 2010.
- [39] R. Connelly. Rigidity. In *Handbook of convex geometry, Vol. A, B*, pages 223–271. North-Holland, Amsterdam, 1993.
- [40] R. Connelly and J.-M. Schlenker. On the infinitesimal rigidity of weakly convex polyhedra. *European J. Combin.*, 31(4):1080–1090, 2010.

- [41] H. T. Croft, K. J. Falconer, and R. K. Guy. *Unsolved problems in geometry*. Problem Books in Mathematics. Springer-Verlag, New York, 1991. Unsolved Problems in Intuitive Mathematics, II.
- [42] H. T. Croft, K. J. Falconer, and R. K. Guy. *Unsolved problems in geometry*. Springer-Verlag, New York, 1994. Corrected reprint of the 1991 original [MR 92c:52001], Unsolved Problems in Intuitive Mathematics, II.
- [43] B. E. J. Dahlberg. The converse of the four vertex theorem. *Proc. Amer. Math. Soc.*, 133(7):2131–2135 (electronic), 2005.
- [44] E. D. Demaine, M. L. Demaine, V. Hart, J. Iacono, S. Langerman, and J. O’Rourke. Continuous blooming of convex polyhedra. *Graphs Combin.*, 27(3):363–376, 2011.
- [45] E. D. Demaine, J. S. B. Mitchell, and J. O’Rourke. The open problems project, cs.smith.edu/~orourke/topp/.
- [46] E. D. Demaine and J. O’Rourke. *Geometric folding algorithms*. Cambridge University Press, Cambridge, 2007. Linkages, origami, polyhedra.
- [47] D. DeTurck, H. Gluck, D. Pomerleano, and D. S. Vick. The four vertex theorem and its converse. *Notices Amer. Math. Soc.*, 54(2):192–207, 2007.
- [48] S. Donaldson. Some problems in differential geometry and topology. *Nonlinearity*, 21(9):T157, 2008.
- [49] A. Dürer. *The painter’s manual: A manual of measurement of lines, areas, and solids by means of compass and ruler assembled by Albrecht Dürer for the use of all lovers of art with appropriate illustrations arranged to be printed in the year MDXXV*. Abaris Books, New York, N.Y., 1977 (1525).
- [50] R. Earp, F. Brito, W. H. Meeks III, and H. Rosenberg. Structure theorems for constant mean curvature surfaces bounded by a planar curve. *Indiana Univ. Math. J.*, 40(1):333–343, 1991.
- [51] E. Egerváry. On the smallest convex cover of a simple arc of space-curve. *Publ. Math. Debrecen*, 1:65–70, 1949.
- [52] T. Ekholm, B. White, and D. Wienholtz. Embeddedness of minimal surfaces with total boundary curvature at most 4π . *Ann. of Math. (2)*, 155(1):209–234, 2002.
- [53] Y. Eliashberg and N. Mishachev. *Introduction to the h-principle*, volume 48 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2002.
- [54] L. Euler. *Leonhardi Euleri opera postuma mathematica et physica. Tomus prior, tomus alter*. Discovered in 1844 in the St. Petersburg Academy of Sciences. Edited by P. H. Fuss and N. Fuss. Kraus Reprint, Nedeln/Liechtenstein, 1969.
- [55] E. A. Feldman. On parabolic and umbilic points of immersed hypersurfaces. *Trans. Amer. Math. Soc.*, 127:1–28, 1967.
- [56] S. R. Finch and J. E. Wetzel. Lost in a forest. *Amer. Math. Monthly*, 111(8):645–654, 2004.
- [57] R. Garcia and C. Gutierrez. Ovaloids of \mathbf{R}^3 and their umbilics: a differential equation approach. *J. Differential Equations*, 168(1):200–211, 2000. Special issue in celebration of Jack K. Hale’s 70th birthday, Part 1 (Atlanta, GA/Lisbon, 1998).
- [58] R. J. Gardner. *Geometric tomography*. Cambridge University Press, Cambridge, 1995.
- [59] C. F. Gauss. *General investigations of curved surfaces*. Translated from the Latin and German by Adam Hildebeitel and James Morehead. Raven Press, Hewlett, N.Y., 1965.
- [60] M. Ghomi. Solution to the shadow problem in 3-space. *Minimal Surfaces, Geometric Analysis and Symplectic Geometry*, pages 129–142, 2000.
- [61] M. Ghomi. Strictly convex submanifolds and hypersurfaces of positive curvature. *J. Differential Geom.*, 57(2):239–271, 2001.
- [62] M. Ghomi. Gauss map, topology, and convexity of hypersurfaces with nonvanishing curvature. *Topology*, 41(1):107–117, 2002.
- [63] M. Ghomi. Shadows and convexity of surfaces. *Ann. of Math. (2)*, 155(1):281–293, 2002.
- [64] M. Ghomi. Classical open problems in differential geometry. www.math.gatech.edu/~ghomi/Classes/Math497C/op.pdf, 2004.
- [65] M. Ghomi. Topology of surfaces with connected shades. *Asian J. Math.*, 11(4):621–634, 2007.

- [66] M. Ghomi. Directed immersions of closed manifolds. *Geom. Topol.*, 15(2):699–705, 2011.
- [67] M. Ghomi. A Riemannian four vertex theorem for surfaces with boundary. *Proc. Amer. Math. Soc.*, 139(1):293–303, 2011.
- [68] M. Ghomi. Tangent lines, inflections, and vertices of closed curves. *Duke Math. J.*, 162(14):2691–2730, 2013.
- [69] M. Ghomi. Vertices of closed curves in Riemannian surfaces. *Comment. Math. Helv.*, 88(2):427–448, 2013.
- [70] M. Ghomi. Affine unfoldings of convex polyhedra. *Geom. Topol.*, 18(5):3055–3090, 2014.
- [71] M. Ghomi. The length, width, and inradius of space curves. *arXiv:1605.01144*, 2016.
- [72] M. Ghomi. Boundary torsion and convex caps of locally convex surfaces. *J. Differential Geom.*, To appear.
- [73] M. Ghomi and R. Howard. Normal curvatures of asymptotically constant graphs and Carathéodory’s conjecture. *Proc. Amer. Math. Soc.*, 140(12):4323–4335, 2012.
- [74] M. Ghomi and M. Kossowski. h -principles for hypersurfaces with prescribed principal curvatures and directions. *Trans. Amer. Math. Soc.*, 358(10):4379–4393 (electronic), 2006.
- [75] É. Ghys, S. Tabachnikov, and V. Timorin. Osculating curves: around the Tait-Kneser theorem. *Math. Intelligencer*, 35(1):61–66, 2013.
- [76] H. Gluck. Almost all simply connected closed surfaces are rigid. In *Geometric topology (Proc. Conf., Park City, Utah, 1974)*, pages 225–239. Lecture Notes in Math., Vol. 438. Springer, Berlin, 1975.
- [77] H. Gluck and L.-H. Pan. Embedding and knotting of positive curvature surfaces in 3-space. *Topology*, 37(4):851–873, 1998.
- [78] J. Graver, B. Servatius, and H. Servatius. *Combinatorial rigidity*, volume 2 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1993.
- [79] R. E. Greene and H. Wu. On the rigidity of punctured ovaloids. *Ann. of Math. (2)*, 94:1–20, 1971.
- [80] R. E. Greene and H. Wu. On the rigidity of punctured ovaloids. II. *J. Differential Geometry*, 6:459–472, 1972. Collection of articles dedicated to S. S. Chern and D. C. Spencer on their sixtieth birthdays.
- [81] M. Gromov. *Partial differential relations*. Springer-Verlag, Berlin, 1986.
- [82] M. Gromov. Spaces and questions. *Geom. Funct. Anal.*, (Special Volume, Part I):118–161, 2000. GAFA 2000 (Tel Aviv, 1999).
- [83] M. Gromov. Number of questions. www.ihes.fr/~gromov/PDF/problems-sept2014-copy.pdf, 2014.
- [84] M. Gromov. Geometric, algebraic and analytic descendants of Nash isometric embedding theorems. www.ihes.fr/~gromov/topics/recent.html, 2015.
- [85] B. Grünbaum. Nets of polyhedra. II. *Geombinatorics*, 1(3):5–10, 1991.
- [86] B. Grünbaum. No-net polyhedra. *Geombinatorics*, 11(4):111–114, 2002.
- [87] B. Guan and J. Spruck. Locally convex hypersurfaces of constant curvature with boundary. *Comm. Pure Appl. Math.*, 57(10):1311–1331, 2004.
- [88] B. Guilfoyle and W. Klingenberg. Proof of the Carathéodory conjecture by mean curvature flow in the space of oriented affine lines. *arXiv:0808.0851v1*, 2008.
- [89] C. Gutierrez and J. Sotomayor. Lines of curvature, umbilic points and Carathéodory conjecture. *Resenhas*, 3(3):291–322, 1998.
- [90] J. Hadamard. Les surfaces a courbures opposees et leurs lignes geodesiques. *J. Math. Pures Appl.*, 4(1):27–73, 1898.
- [91] H. Hamburger. Beweis einer Carathéodoryschen Vermutung. Teil I. *Ann. of Math. (2)*, 41:63–86, 1940.
- [92] H. L. Hamburger. Beweis einer Carathéodoryschen Vermutung. II. *Acta Math.*, 73:175–228, 1941.
- [93] H. L. Hamburger. Beweis einer Carathéodoryschen Vermutung. III. *Acta Math.*, 73:229–332, 1941.

- [94] Q. Han and J.-X. Hong. *Isometric embedding of Riemannian manifolds in Euclidean spaces*, volume 130 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2006.
- [95] E. M. Harrell, II. A direct proof of a theorem of Blaschke and Lebesgue. *J. Geom. Anal.*, 12(1):81–88, 2002.
- [96] J.-B. Hiriart-Urruty. Du calcul différentiel au calcul variationnel: un aperçu de l'évolution de p. fermata nos jours. *Quadrature*, (70):8–18, 2008.
- [97] D. Hoffman and W. H. Meeks. Minimal surfaces based on the catenoid. *American Mathematical Monthly*, 97(8):702–730, 1990.
- [98] B. K. P. Horn and M. J. Brooks, editors. *Shape from shading*. MIT Press, Cambridge, MA, 1989.
- [99] J.-i. Itoh, J. O'Rourke, and C. Vilcu. Star unfolding convex polyhedra via quasigeodesic loops. *Discrete Comput. Geom.*, 44(1):35–54, 2010.
- [100] P. Ivanisvili. Burkholder's Martingale transform. *arXiv:1402.4751*, 2014.
- [101] P. Ivanisvili, D. M. Stolyarov, and P. B. Zatitskiy. Bellman VS Beurling: sharp estimates of uniform convexity for L^p spaces. *arXiv:1405.6229*, 2014.
- [102] V. V. Ivanov. An analytic conjecture of Carathéodory. *Sibirsk. Mat. Zh.*, 43(2):314–405, ii, 2002.
- [103] I. Izmestiev and J.-M. Schlenker. Infinitesimal rigidity of polyhedra with vertices in convex position. *Pacific J. Math.*, 248(1):171–190, 2010.
- [104] W. C. Jagy. Minimal hypersurfaces foliated by spheres. *Michigan Math. J.*, 38(2):255–270, 1991.
- [105] L. P. d. M. Jorge and F. Xavier. A complete minimal surface in \mathbb{R}^3 between two parallel planes. *Annals of Mathematics*, pages 203–206, 1980.
- [106] N. Kapouleas et al. Compact constant mean curvature surfaces in euclidean three-space. *Journal of Differential Geometry*, 33(3):683–715, 1991.
- [107] R. Kirby. Problems in low-dimensional topology. www.math.berkeley.edu/~kirby, 1995.
- [108] Y. Kitagawa. Deformable flat tori in S^3 with constant mean curvature. *Osaka J. Math.*, 40(1):103–119, 2003.
- [109] F. Klein. Eine neue Relation zwischen den Singularitäten einer algebraischen Curve. *Math. Ann.*, 10(2):199–209, 1876.
- [110] T. Klotz. On G. Bol's proof of Carathéodory's conjecture. *Comm. Pure Appl. Math.*, 12:277–311, 1959.
- [111] A. Kneser. Bemerkungen über die anzahl der extrema des krümmung auf geschlossenen kurven und über verwandte fragen in einer nicht euklidischen geometrie. In *Festschrift Heinrich Weber*, pages 170–180. Teubner, 1912.
- [112] G. A. Kovalëva. Absence of closed asymptotic lines on tubes of negative Gaussian curvature with a one-to-one spherical mapping. *Fundam. Prikl. Mat.*, 1(4):953–977, 1995.
- [113] N. H. Kuiper. Geometry in curvature theory. In *Tight and taut submanifolds (Berkeley, CA, 1994)*, pages 1–50. Cambridge Univ. Press, Cambridge, 1997.
- [114] L. Lazarovici. Elliptic sectors in surface theory and the Carathéodory-Loewner conjectures. *J. Differential Geom.*, 55(3):453–473, 2000.
- [115] L. Liberti and C. Lavor. Six mathematical gems from the history of distance geometry. *arXiv:1502.02816*, 2015.
- [116] F. López, F. Martín, S. Morales, et al. Adding handles to nadirashvili's surfaces. *J. Differential Geom.*, 60(1):155–175, 2002.
- [117] R. López. *Constant mean curvature surfaces with boundary*. Springer Science & Business Media, 2013.
- [118] B. Lucier. Local overlaps in special unfoldings of convex polyhedra. *Comput. Geom.*, 42(5):495–504, 2009.
- [119] E. Makai, Jr. On the geodesic diameter of convex surfaces. *Period. Math. Hungar.*, 4:157–161, 1973.

- [120] N. P. Makuha. A connection between the inner and outer diameters of a general closed convex surface. *Ukrain. Geometr. Sb. Vyp.*, 2:49–51, 1966.
- [121] N. P. Makuha. An isoperimetric property for surfaces of rotation. *Ukrain. Geometr. Sb. Vyp.*, 3:55–56, 1966.
- [122] F. C. Marques and A. Neves. Min-max theory and the Willmore conjecture. *Ann. of Math. (2)*, 179(2):683–782, 2014.
- [123] F. Martín, S. Morales, et al. Complete proper minimal surfaces in convex bodies of \mathbb{R}^n . *Duke Mathematical Journal*, 128(3):559–593, 2005.
- [124] J. Maxwell. On the transformation of surfaces by bending. *Tran. Camb. Phil. Soc.*, 9(4):445–470, 1819.
- [125] W. Meeks. *Lectures on Plateau’s problem*. UFCE, 1978.
- [126] W. H. Meeks and M. Weber. Minimal surfaces. *Oberwolfach Reports*, 6(4):2545–2584, 2009.
- [127] W. H. Meeks, III and B. White. Minimal surfaces bounded by convex curves in parallel planes. *Comment. Math. Helv.*, 66(2):263–278, 1991.
- [128] W. H. Meeks, III and B. White. The space of minimal annuli bounded by an extremal pair of planar curves. *Comm. Anal. Geom.*, 1(3-4):415–437, 1993.
- [129] W. H. Meeks III. Global problems in classical minimal surface theory. *Global theory of minimal surfaces*, pages 453–470, 2005.
- [130] Z. A. Melzak. The isoperimetric problem of the convex hull of a closed space curve. *Proc. Amer. Math. Soc.*, 11:265–274, 1960.
- [131] E. Miller and I. Pak. Metric combinatorics of convex polyhedra: cut loci and nonoverlapping unfoldings. *Discrete Comput. Geom.*, 39(1-3):339–388, 2008.
- [132] T. K. Milnor. Efimov’s theorem about complete immersed surfaces of negative curvature. *Advances in Math.*, 8:474–543, 1972.
- [133] A. F. Möbius. Über die grundformen der linien der dritten ordnung. In *Gesammelte Werke II*, pages 89–176. Verlag von S. Hirzel, Leipzig, 1886.
- [134] H. Mohrmann. Die minimalzahl der stationären ebenen eines räumlichen ovals. *Sitz Ber Kgl Bayerischen Akad. Wiss. Math. Phys. Kl.*, pages 1–3, 1917.
- [135] S. Mukhopadhyaya. New methods in the geometry of a plane arc. *Bull. Calcutta Math. Soc. I*, pages 31–37, 1909.
- [136] N. Nadirashvili. Hadamard’s and calabi–yau’s conjectures on negatively curved and minimal surfaces. *Inventiones mathematicae*, 126(3):457–465, 1996.
- [137] N. Nadirashvili and Y. Yuan. Improving Pogorelov’s isometric embedding counterexample. *Calc. Var. Partial Differential Equations*, 32(3):319–323, 2008.
- [138] B. Nelli. An example of an immersed complete genus one minimal surface in \mathbb{R}^3 with two convex ends. *Comment. Math. Helv.*, 73(2):298–305, 1998.
- [139] I. Nikolaev. *Foliations on surfaces*, volume 41 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2001. With a foreword by Idel Bronshteyn and Chapter 16 by B. Piccoli.
- [140] L. Nirenberg. Rigidity of a class of closed surfaces. In *Nonlinear Problems (Proc. Sympos., Madison, Wis., 1962)*, pages 177–193. Univ. of Wisconsin Press, Madison, Wis., 1963.
- [141] J. J. Nuño Ballesteros and M. C. Romero Fuster. A four vertex theorem for strictly convex space curves. *J. Geom.*, 46(1-2):119–126, 1993.
- [142] J. O’Rourke. *How to fold it*. Cambridge University Press, Cambridge, 2011. The mathematics of linkages, origami, and polyhedra.
- [143] J. O’Rourke. Shortest closed curve to inspect a sphere (question posted on mathoverflow). mathoverflow.net/questions/69099/shortest-closed-curve-to-inspect-a-sphere, June 2011.
- [144] V. Ovsienko and S. Tabachnikov. *Projective differential geometry old and new*, volume 165 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2005. From the Schwarzian derivative to the cohomology of diffeomorphism groups.
- [145] I. Pak. *Lectures on discrete and polyhedral geometry*. www.math.umn.edu/~pak/, 2008.

- [146] J. Pardon. On the distortion of knots on embedded surfaces. *Ann. Math.*, 174(1):637–646, 2011.
- [147] U. Pinkall. Hopf tori in S^3 . *Invent. Math.*, 81(2):379–386, 1985.
- [148] A. V. Pogorelov. An example of a two-dimensional Riemannian metric that does not admit a local realization in E_3 . *Dokl. Akad. Nauk SSSR*, 198:42–43, 1971.
- [149] A. V. Pogorelov. *Extrinsic geometry of convex surfaces*. American Mathematical Society, Providence, R.I., 1973. Translated from the Russian by Israel Program for Scientific Translations, Translations of Mathematical Monographs, Vol. 35.
- [150] E. Raphael, J.-M. di Meglio, M. Berger, and E. Calabi. Convex particles at interfaces. *J. Phys. I France*, 2:571–579, 1992.
- [151] E. Rembs. Zur Verbiegung von Flächen im Grossen. *Math. Z.*, 56:271–279, 1952.
- [152] M. Rochowski. Special problems of surface theory in the Euclidean 3-dimensional space. *Demonstratio Math.*, 25(1-2):1–23, 1992.
- [153] L. Rodríguez and H. Rosenberg. Rigidity of certain polyhedra in \mathbf{R}^3 . *Comment. Math. Helv.*, 75(3):478–503, 2000.
- [154] M. C. Romero-Fuster and V. D. Sedykh. A lower estimate for the number of zero-torsion points of a space curve. *Beiträge Algebra Geom.*, 38(1):183–192, 1997.
- [155] A. Ros. Embedded minimal surfaces: forces, topology and symmetries. *Calc. Var. Partial Differential Equations*, 4(5):469–496, 1996.
- [156] A. Ros and H. Rosenberg. Constant mean curvature surfaces in a half-space of \mathbf{r}^3 with boundary in the boundary of the half-space. *J. Differential Geom.*, 44(4):807–817, 1996.
- [157] H. Rosenberg. Hypersurfaces of constant curvature in space forms. *Bull. Sci. Math.*, 117(2):211–239, 1993.
- [158] C. Rourke and B. Sanderson. The compression theorem. II. Directed embeddings. *Geom. Topol.*, 5:431–440 (electronic), 2001.
- [159] È. R. Rozendorn. Surfaces of negative curvature [MR1039819 (92a:53087)]. In *Geometry, III*, volume 48 of *Encyclopaedia Math. Sci.*, pages 87–178, 251–256. Springer, Berlin, 1992.
- [160] R. Sacksteder. On hypersurfaces with no negative sectional curvatures. *Amer. J. Math.*, 82:609–630, 1960.
- [161] T. Sakai. On the isodiametric inequality for the 2-sphere. In *Geometry of manifolds (Matsumoto, 1988)*, volume 8 of *Perspect. Math.*, pages 303–315. Academic Press, Boston, MA, 1989.
- [162] H. Scherbel. *A new proof of Hamburger’s Index Theorem on umbilical points*. ProQuest LLC, Ann Arbor, MI, 1993. Thesis (Dr.Sc.Math)–Eidgenoessische Technische Hochschule Zuerich (Switzerland).
- [163] C. Schevon. *Algorithms for geodesics on polytopes*. PhD thesis, Johns Hopkins University, 1989.
- [164] W. Schlickerieder. Nets of polyhedra. Master’s thesis, Technische Universität Berlin, 1997.
- [165] R. Schneider. *Convex bodies: the Brunn-Minkowski theory*. Cambridge University Press, Cambridge, 1993.
- [166] R. Schoen and S.-T. Yau. *Lectures on differential geometry*. Conference Proceedings and Lecture Notes in Geometry and Topology, I. International Press, Cambridge, MA, 1994. Lecture notes prepared by Wei Yue Ding, Kung Ching Chang [Gong Qing Zhang], Jia Qing Zhong and Yi Chao Xu, Translated from the Chinese by Ding and S. Y. Cheng, Preface translated from the Chinese by Kaising Tso.
- [167] R. M. Schoen. Uniqueness, symmetry, and embeddedness of minimal surfaces. *J. Differential Geom.*, 18(4):791–809 (1984), 1983.
- [168] I. J. Schoenberg. An isoperimetric inequality for closed curves convex in even-dimensional Euclidean spaces. *Acta Math.*, 91:143–164, 1954.
- [169] V. D. Sedykh. Four vertices of a convex space curve. *Bull. London Math. Soc.*, 26(2):177–180, 1994.

- [170] G. C. Shephard. Convex polytopes with convex nets. *Math. Proc. Cambridge Philos. Soc.*, 78(3):389–403, 1975.
- [171] M. Shiffman. On surfaces of stationary area bounded by two circles, or convex curves, in parallel planes. *Ann. of Math. (2)*, 63:77–90, 1956.
- [172] T. Shioya. Diameter and area estimates for S^2 and \mathbf{P}^2 with non-negatively curved metrics. In *Progress in differential geometry*, volume 22 of *Adv. Stud. Pure Math.*, pages 309–319. Math. Soc. Japan, Tokyo, 1993.
- [173] B. Smyth and F. Xavier. A sharp geometric estimate for the index of an umbilic on a smooth surface. *Bull. London Math. Soc.*, 24(2):176–180, 1992.
- [174] B. Smyth and F. Xavier. Real solvability of the equation $\partial_{\bar{z}}^2\omega = \rho g$ and the topology of isolated umbilics. *J. Geom. Anal.*, 8(4):655–671, 1998.
- [175] B. Smyth and F. Xavier. Eigenvalue estimates and the index of Hessian fields. *Bull. London Math. Soc.*, 33(1):109–112, 2001.
- [176] J. Sotomayor and R. Garcia. Lines of curvature on surfaces, historical comments and recent developments. *São Paulo J. Math. Sci.*, 2(1):99–143, 2008.
- [177] D. Spring. Directed embeddings of closed manifolds. *Commun. Contemp. Math.*, 7(5):707–725, 2005.
- [178] J. J. Stoker. Geometrical problems concerning polyhedra in the large. *Comm. Pure Appl. Math.*, 21:119–168, 1968.
- [179] D. J. Struik. Differential geometry in the large. *Bull. Amer. Math. Soc.*, 37(2):49–62, 1931.
- [180] A. tarasov. Existence of a polyhedron which does not have a non-overlapping pseudo-edge unfolding. *arXiv:0806.2360v3*, 2008.
- [181] G. Thorbergsson and M. Umehara. A unified approach to the four vertex theorems. II. In *Differential and symplectic topology of knots and curves*, volume 190 of *Amer. Math. Soc. Transl. Ser. 2*, pages 229–252. Amer. Math. Soc., Providence, RI, 1999.
- [182] C. J. Titus. A proof of a conjecture of Loewner and of the conjecture of Caratheodory on umbilic points. *Acta Math.*, 131:43–77, 1973.
- [183] V. A. Toponogov. On conditions for the existence of umbilical points on a convex surface. *Sibirsk. Mat. Zh.*, 36(4):903–910, iv, 1995.
- [184] N. S. Trudinger and X.-J. Wang. On locally convex hypersurfaces with boundary. *J. Reine Angew. Math.*, 551:11–32, 2002.
- [185] J. Van Heijenoort. On locally convex manifolds. *Comm. Pure Appl. Math.*, 5:223–242, 1952.
- [186] F. J. Wegner. Floating bodies of equilibrium in 2d, the tire track problem and electrons in a parabolic magnetic field. *arXiv preprint physics/0701241*, 2007.
- [187] H. Wente. Personal email. January 9, 1999.
- [188] S. T. Yau. Problem section. In *Seminar on Differential Geometry*, volume 102 of *Ann. of Math. Stud.*, pages 669–706. Princeton Univ. Press, Princeton, N.J., 1982.
- [189] S.-T. Yau. Open problems in geometry. In *Differential geometry: partial differential equations on manifolds (Los Angeles, CA, 1990)*, pages 1–28. Amer. Math. Soc., Providence, RI, 1993.
- [190] S.-T. Yau. Review of geometry and analysis. In *Mathematics: frontiers and perspectives*, pages 353–401. Amer. Math. Soc., Providence, RI, 2000.
- [191] V. A. Zalgaller, editor. *Geometry. III*, volume 48 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 1992. Theory of surfaces, A translation of *Current problems in mathematics. Fundamental directions. Vol. 48* (Russian), Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1989 [MR1039817 (90j:53002)], Translation by E. Primrose, Translation edited by Yu. D. Burago and V. A. Zalgaller.
- [192] V. A. Zalgaller. The problem of the shortest space curve of unit width. *Mat. Fiz. Anal. Geom.*, 1(3-4):454–461, 1994.
- [193] V. A. Zalgaller. Extremal problems on the convex hull of a space curve. *Algebra i Analiz*, 8(3):1–13, 1996.
- [194] V. A. Zalgaller. Shortest inspection curves for a sphere. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)*, 299(Geom. i Topol. 8):87–108, 328, 2003.

- [195] G. M. Ziegler. *Lectures on polytopes*, volume 152 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332

E-mail address: ghomi@math.gatech.edu

URL: www.math.gatech.edu/~ghomi