

Quiz 3 for Calculus ++, Math 2605A1-2, October 12, 2004

Name:

This quiz is to be taken without calculators and notes of any sorts. The allowed time is 20 minutes. Provide exact answers; not decimal approximations! For example, if you mean $\sqrt{2}$ do not write 1.414...

I: Consider the matrix

$$A = \begin{bmatrix} 2 & 1 & 3 \\ 1 & 6 & 1 \\ 3 & 1 & 2 \end{bmatrix}.$$

a) (3 points) Using the largest off-diagonal elements for the first step in the Jacobi Algorithm write down the Givens rotation G and calculate the matrix $A^{(1)} = G^T A G$.

b) (1 point) Find an exact eigenvalue of the matrix A .

II: (3 points) Recall that $\text{Off}(A)$ denotes the sum of the squares of the off-diagonal elements of A and $A^{(k)}$ is the matrix after the k -th step in the Jacobi algorithm. Which of the statements are true and which are false:

a) $\text{Off}(A^{(12)}) \geq \text{Off}(A^{(13)})$

b) $\text{Off}(A^{(k)}) \geq \frac{1}{1+k^2}$

c) $\text{Off}(A^{(13)}) \leq \frac{2}{3} \text{Off}(A^{(12)})$

III: (3 points) Consider the matrix

$$\begin{bmatrix} 3 & 1 & 0 \\ 1 & 3 & 0 \\ 0 & 0 & 5 \end{bmatrix} + t \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 3 & 4 & 5 \end{bmatrix}.$$

Find all the eigenvalues for small values of t , i.e., in the forms $\mu_i(0) + \mu'_i(0)t + o(t)$, $i = 1, 2, 3$ with explicit values for $\mu_i(0)$ and $\mu'_i(0)$.