



c. Use your answer to part (a) to factor 204 into primes. You will need this answer later in the test.

4. For a positive integer n , let t_n count the number of ternary sequences which do not contain 2011 as four consecutive characters. Develop a recurrence for t_n and use it to find t_6 .


5. Use the algorithm developed in class (from node i , always take the edge ij not traversed previously, where j is minimum) to find an Euler circuit in the graph G shown below. Use node 1 as root.



6. Show that the graph below is hamiltonian. You may give your answer by darkening appropriate edges on the figure, or by giving an appropriate permutation of the vertex set.




7. Consider the graph G shown below.



- For this graph, $\omega(G)$ is:
- Show that $\chi(G) = \omega(G)$ by providing a proper coloring of G . You may indicate your coloring by writing directly on the figure.
- Explain why the graph G is *not* perfect.

8. Consider the poset P shown below.



- The poset P is not an interval order. By inspection, find four points which determine a subposet isomorphic to $\mathbf{2} + \mathbf{2}$:
- List a set of elements which forms a maximum antichain in P :

c. By inspection, find a Dilworth partition of the poset P . You may provide your answer by writing directly on the figure.

9. For the subset lattice 2^9 ,


a. The total number of elements is:

b. The total number of maximal chains is:

c. The number of maximal chains through $\{3, 5, 8\}$ is:

d. The width of 2^9 is:


10. Consider the poset P shown below.



a. Determine a minimum partition of P into antichains by recursively stripping off the set of minimal elements. You may display your answer by writing directly on the diagram. Then darken a set of points that form a maximum chain.

b. The height h of the poset P is:


11. The poset P shown below is an interval order:



a. On the next page, find the down sets and the up sets. Then use these answers to find an interval representation of P that uses the least number of end points.

$D(a) =$	$U(a) =$	$I(a) =$
$D(b) =$	$U(b) =$	$I(b) =$
$D(c) =$	$U(c) =$	$I(c) =$
$D(d) =$	$U(d) =$	$I(d) =$
$D(e) =$	$U(e) =$	$I(e) =$
$D(f) =$	$U(f) =$	$I(f) =$
$D(g) =$	$U(g) =$	$I(g) =$
$D(h) =$	$U(h) =$	$I(h) =$

b. In the space below, draw the representation you have found. Then use the First Fit Coloring Algorithm for interval graphs to solve the Dilworth Problem for this poset, i.e., find the width w and a partition of P into w chains. You may display your answers by writing the colors directly on the intervals in the diagram.



c. The following points form a maximum antichain in P :

12 a. Write all the partitions of the integer 5.

b. Returning to your answer to part a, mark all the partitions of 5 into odd parts with a capital O . Also mark all partitions of 5 into distinct parts with a capital D .

13a. Find the general solution to the advancement operator equation:

$$A^4(A - 5 + 4i)^3(A - 1)^2(A + 8)(A - 9)f = 0$$

b. Write the form of a particular solution of the non-homogeneous advancement operator equation (do not carry out the work necessary to evaluate any constants in your answer):

$$A^4(A - 5 + 4i)^3(A - 1)^2(A + 8)(A - 9)f = 17 \cdot 3^n - 28 \cdot 2^n.$$

c. Find the solution to the advancement operator equation:

$$(A^2 - 11A + 28)f(n) = 0, \quad f(0) = -2 \text{ and } f(1) = 1.$$

14a. Write the inclusion/exclusion formula for the number $S(n, m)$ of onto functions from $\{1, 2, \dots, n\}$ to $\{1, 2, \dots, m\}$.


b. Evaluate your answer to part a when $n = 5$ and $m = 3$ (reduce your answer so that only arithmetic is required to obtain the final answer).

c. Write the inclusion/exclusion formula for the number d_n of derangements on $\{1, 2, \dots, n\}$.


c. Evaluate your formula for d_n when $n = 7$ (reduce your answer so that only arithmetic is required to get the final answer).

d. Use your answer to Problem 3(c) to find the value of the Euler ϕ -function $\phi(n)$ when $n = 204$ (reduce your answer so that only arithmetic is required to get the final answer).

15 In the space to the right, verify Euler's formula for the following planar graph:



16 Consider the following weighted graph:



In the space below, list *in order* the edges which make up a minimum weight spanning tree using, respectively Kruskal's Algorithm (avoid cycles) and Prim's Algorithm (build tree). For Prim, use vertex *A* as the root.

Kruskal's Algorithm

Prim's Algorithm

17. Consider a poset P whose ground set is $X = \{1, 2, 3, \dots, 9\}$. Network flows (and the special case of bipartite matchings) are used to find the width w of P and a minimum chain partition. When the labelling algorithm halts, the following edges are matched:

$$3'8'' \quad 4'6'' \quad 5'7'' \quad 2'1'' \quad 7'2''$$


a. Find the chain partition of P that is associated with this matching.

b. Find the width w of the poset P .

18. A data file `digraph_data.txt` has been read for a digraph whose vertex set is $[7]$. The weights on the directed edges are shown in the matrix below. The entry $w(i, j)$ denotes the length of the edge from i to j . If there is no entry, then the edge is not present in the graph. Apply Dijkstra's algorithm to find the distance $d(x)$ from vertex 1 to vertex x for all vertices x in G . Also, for each x , find a shortest path $P(x)$ from 1 to x .

W	1	2	3	4	5	6	7
1	0	20	17	28	23	44	48
2		0	4	6	3	8	16
3		9	0		41	12	20
4	27	20	28	0		2	10
5	9	1	2	2	0	4	9
6	82	5	3	2	2	0	7
7		8	22	4	18	12	0

19. Consider the following network flow:



- a. The value of the current flow is:
- b. The capacity of the cut $\{S, B, D, G\} \cup \{A, C, E, F, T\}$ is:
- c. Carry out the labeling algorithm, using the pseudo-alphabetic order on the vertices and list below the labels which will be given to the vertices.
- d. Use your work in part c to find an augmenting path and make the appropriate changes directly on the diagram.
- e. Carry out the labeling algorithm a second time on the updated flow. It should halt without the sink being labeled.
- f. Find a cut whose capacity is equal to the value of the updated flow.

20. True–False. Mark in the left margin.

1. $2^{40} > 100,000,000$.
2. The coefficient of $x^6y^8z^2$ in $(2x - 4y^3 + 8z)^{29}$ is 0.
3. There is a planar graph G on 328 vertices with $\chi(G) = 39$.
4. If G is a planar graph, then $\chi(G) = \omega(G)$.
5. All graphs with 1024 vertices and 2973 edges are planar.
6. Every graph on 684 vertices in which every vertex has degree 402 is hamiltonian.
7. Every connected graph on 986 vertices in which every vertex has degree 42 has an Euler circuit.
8. A cycle on 739 vertices is a homeomorph of the complete bipartite graph $K_{2,2}$.
9. When $n \geq 3$, the shift graph S_n is a triangle-free graph with $\binom{n}{2}$ vertices and $\binom{n}{3}$ edges. Furthermore, $\chi(S_n) = \lceil \lg n \rceil$.
10. The number of lattice paths from $(0, 0)$ to (n, n) which do not pass through a point above the diagonal is the Catalan number $\binom{2n}{n}/(n+1)$.
11. Any modern computer can accept a file of 1,000 positive integers, each at most 5,000, and quickly determine whether 2,742 is the sum of three integers in the file.
12. Any modern computer can accept a file of 1,000 positive integers, each at most 5,000, and quickly determine whether 385,742 is the product of two integers in the file.
13. Any modern computer can accept a file of 1,000 positive integers, each at most 5,000, and quickly factor each of the numbers into primes.
14. There is a graph on 837 vertices in which no two vertices have the same degree.
15. There is a poset with 623 points having width 29 and height 19.
16. There is a sequence of 523 distinct positive integers which does not have an increasing subsequence of size 38 nor a decreasing subsequence of size 27.
17. The permutation $(7, 1, 3, 8, 5, 2, 4, 6)$ is a derangement.
18. Linear programming problems with integer coefficient constraints always have integer valued solutions.
19. Every linear programming problem is also a network flow problem.
20. The Ramsey number $R(3, 3)$ is 18.