



An Ansatz for the asymptotics of hypergeometric multisums

Stavros Garoufalidis¹

School of Mathematics, Georgia Institute of Technology, Atlanta, GA 30332-0160, USA

Received 12 July 2007; accepted 12 February 2008

Available online 31 July 2008

Abstract

Sequences that are defined by multisums of hypergeometric terms with compact support occur frequently in enumeration problems of combinatorics, algebraic geometry and perturbative quantum field theory. The standard recipe to study the asymptotic expansion of such sequences is to find a recurrence satisfied by them, convert it into a differential equation satisfied by their generating series, and analyze the singularities in the complex plane. We propose a shortcut by constructing directly from the structure of the hypergeometric term a finite set, for which we conjecture (and in some cases prove) that it contains all the singularities of the generating series. Our construction of this finite set is given by the solution set of a balanced system of polynomial equations of a rather special form, reminiscent of the Bethe ansatz. The finite set can also be identified with the set of critical values of a potential function, as well as with the evaluation of elements of an additive K -theory group by a regulator function. We give a proof of our conjecture in some special cases, and we illustrate our results with numerous examples.

© 2008 Elsevier Inc. All rights reserved.

MSC: primary 57N10; secondary 57M25

Keywords: Holonomic functions; Regular holonomic functions; Balanced hypergeometric terms; WZ algorithm; Diagonals of rational functions; Asymptotic expansions; Transseries; Laplace transform; Hadamard product; Bethe ansatz; Singularities; Additive K -theory; Regulators; Stirling formula; Algebraic combinatorics; Polytopes; Newton polytopes; G -functions

E-mail address: stavros@math.gatech.edu.

URL: <http://www.math.gatech.edu/~stavros>.

¹ The author was supported in part by NSF.

Contents

1.	Introduction	424
1.1.	The problem	424
1.2.	Existence of asymptotic expansions	424
1.3.	Computation of asymptotic expansions	425
1.4.	Hypergeometric terms	426
1.5.	Balanced terms, generating series and singularities	427
1.6.	The definition of S_t and K_t	428
1.7.	The conjecture	429
1.8.	Partial results	429
1.9.	Laurent polynomials: A source of special terms	430
1.10.	Plan of the proof	431
2.	Balanced terms and potential functions	431
2.1.	The Stirling formula and potential functions	431
2.2.	Proof of Theorem 5	433
3.	Balanced terms, the entropy function and additive K -theory	434
3.1.	A brief review of the entropy function and additive K -theory	434
3.2.	Balanced terms and additive K -theory	436
3.3.	Proof of Theorem 6	437
4.	Proof of Theorems 2 and 3	438
4.1.	Proof of Theorem 2	438
4.2.	Proof of Theorem 3	439
5.	Some examples	440
5.1.	A closed form example	440
5.2.	The Apery sequence	441
5.3.	An example with critical points at the boundary	442
6.	Laurent polynomials and special terms	443
6.1.	Proof of Theorem 4	443
6.2.	The Newton polytope of a Laurent polynomial and its associated balanced term	446
	Acknowledgments	450
	References	450

1. Introduction

1.1. The problem

The problem considered here is the following: given a balanced hypergeometric term t_{n,k_1,\dots,k_r} with compact support for each $n \in \mathbb{N}$, we wish to find an asymptotic expansion of the sequence

$$a_{t,n} = \sum_{(k_1,\dots,k_r) \in \mathbb{Z}^r} t_{n,k_1,\dots,k_r}. \quad (1)$$

Such sequences occur frequently in enumeration problems of combinatorics, algebraic geometry and perturbative quantum field theory; see [10,20,31]. The standard recipe for this is to find a recurrence satisfied by (a_n) , convert it into a differential equation satisfied by the generating series

$$G_t(z) = \sum_{n=0}^{\infty} a_{t,n} z^n$$

and analyze the singularities of its analytic continuation in the complex plane. We propose a shortcut by constructing directly from the structure of the hypergeometric term t_{n,k_1,\dots,k_r} a finite set S_t , for which we conjecture (and in some cases prove) that it contains all the singularities of $G(z)$. Our construction of S_t is given by the solution set of a balanced system of polynomial equations of a rather special form, reminiscent of the Bethe ansatz. S_t can also be identified with the set of critical values of a potential function, as well as with the evaluation of elements of an additive K -theory group by a regulator function. We give a proof of our conjecture in some special cases, and we illustrate our results with numerous examples.

1.2. Existence of asymptotic expansions

A general existence theorem for asymptotic expansions of sequences discussed above was recently given in [14]. To phrase it, we need to recall what is a G -function in the sense of Siegel [30].

Definition 1.1. We say that series $G(z) = \sum_{n=0}^{\infty} a_n z^n$ is a G -function if:

- (a) the coefficients a_n are algebraic numbers, and
- (b) there exists a constant $C > 0$ so that for every $n \in \mathbb{N}$ the absolute value of every conjugate of a_n is less than or equal to C^n , and
- (c) the common denominator of a_0, \dots, a_n is less than or equal to C^n .
- (d) $G(z)$ is holonomic, i.e., it satisfies a linear differential equation with coefficients polynomials in z .

The main result of [14] is the following theorem.

Theorem 1. (See [14, Theorem 3].) For every balanced term t , the generating series $G_t(z)$ is a G -function.

Using the fact that the local monodromy of a G -function around a singular point is *quasi-unipotent* (see [2,3,5,9,18]), an elementary application of Cauchy’s theorem implies the following corollary; see [14,17] and [6, Section 7].

Corollary 1.2. If $G(z) = \sum_{n=0}^{\infty} a_n z^n$ is a G -function, then (a_n) has a transseries expansion, that is an expansion of the form

$$a_n \sim \sum_{\lambda \in \Sigma} \lambda^{-n} n^{\alpha_\lambda} (\log n)^{\beta_\lambda} \sum_{s=0}^{\infty} \frac{c_{\lambda,s}}{n^s} \tag{2}$$

where Σ is the set of singularities of $G(z)$, $\alpha_\lambda \in \mathbb{Q}$, $\beta_\lambda \in \mathbb{N}$, and $c_{\lambda,s} \in \mathbb{C}$. In addition, Σ is a finite set of algebraic numbers, and generates a number field $E = \mathbb{Q}(\Sigma)$.

1.3. Computation of asymptotic expansions

Theorem 1 and its Corollary 1.2 are not constructive. The usual way for computing the asymptotic expansion for sequences $(a_{t,n})$ of the form (1) is to find a linear recurrence, and convert it into a differential equation for the generating series $G_t(z)$. The singularities of $G_t(z)$ are easily located from the roots of coefficient of the leading derivative of the ODE. This approach is taken by Wimp, Zeilberger, following Birkhoff, Trjitzinsky; see [4,34] and also [23]. A recurrence for a multisum sequence $(a_{t,n})$ follows from Wilf–Zeilberger’s constructive theorem, and its computer implementation; see [26–28,33,35]. Although constructive, these algorithms are impractical for multisums with, say, more than three summation variables.

On the other hand, it seems wasteful to compute an ODE for $G_t(z)$, and then discard all but a small part of it in order to determine the singularities Σ_t of $G_t(z)$.

The main result of the paper is a construction of a finite set S_t of algebraic numbers directly from the summand t_{n,k_1,\dots,k_r} , which we conjecture that it includes the set Σ_t . We give a proof of our conjecture in some special cases, as well as supporting examples.

Our definition of the set Σ_t is reminiscent of the Bethe ansatz, and is related to critical values of potential functions and additive K -theory.

Before we formulate our conjecture let us give an instructive example.

Example 1.3. Consider the *Apery sequence* (a_n) defined by

$$a_n = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2. \tag{3}$$

It turns out that (a_n) satisfies a linear recursion relation with coefficients in $\mathbb{Q}[n]$ (see [34, p. 174])

$$(n+2)^3 a_{n+2} - (2n+3)(17n^2 + 51n + 39)a_{n+1} + (n+1)^3 a_n = 0 \tag{4}$$

for all $n \in \mathbb{N}$, with initial conditions $a_0 = 1, a_1 = 5$. It follows that $G(z)$ is *holonomic*; i.e., it satisfies a linear ODE with coefficients in $\mathbb{Q}[z]$:

$$z^2(z^2 - 34z + 1)G'''(z) + 3z(2z^2 - 51z + 1)G''(z) + (7z^2 - 112z + 1)G'(z) + (z - 5)G(z) = 0 \tag{5}$$

with initial conditions $G(0) = 1, G'(0) = 5, G''(0) = 146$.

This implies that the possible singularities of $G(z)$ are the roots of the equation:

$$z^2(z^2 - 34z + 1) = 0. \tag{6}$$

Thus, $G(z)$ has analytic continuation as a multivalued function in $\mathbb{C} \setminus \{0, 17 + 12\sqrt{2}, 17 - 12\sqrt{2}\}$. Since the Taylor series coefficients of $G(z)$ at $z = 0$ are positive integers, and $G(z)$ is analytic at $z = 0$, it follows that $G(z)$ has a singularity inside the punctured unit disk. Thus, $G(z)$ is singular at $17 - 12\sqrt{2}$. By Galois invariance, it is also singular at $17 + 12\sqrt{2}$. The proof of Corollary 1.2 implies that (a_n) has an asymptotic expansion of the form:

$$a_n \sim (17 + 12\sqrt{2})^n n^{-3/2} \sum_{s=0}^{\infty} \frac{c_{1s}}{n^s} + (17 - 12\sqrt{2})^n n^{-3/2} \sum_{s=0}^{\infty} \frac{c_{2s}}{n^s} \tag{7}$$

for some constants $c_{1s}, c_{2s} \in \mathbb{C}$ with $c_{10}c_{20} \neq 0$. A final calculation shows that

$$c_{10} = \frac{1}{\pi^{3/2}} \frac{3 + 2\sqrt{2}}{4\sqrt[4]{2}}, \quad c_{20} = \frac{1}{\pi^{3/2}} \frac{3 - 2\sqrt{2}}{4\sqrt[4]{2}}.$$

Our paper gives an ansatz that quickly produces the numbers $17 \pm 2\sqrt{2}$ given the expression (3) of (a_n) , bypassing Eqs. (4) and (5). This is explained in Section 5.2. In a forthcoming publication we will explain how to compute the *Stokes constants* c_{s0} for $s = 0, 1$ in terms of the expression (3).

1.4. Hypergeometric terms

We have already mentioned multisums of balanced hypergeometric terms. Let us define what those are.

Definition 1.4. An r -dimensional *balanced hypergeometric term* $t_{n,k}$ (in short, *balanced term*, also denoted by t) in variables (n, k) , where $n \in \mathbb{N}$ and $k = (k_1, \dots, k_r) \in \mathbb{Z}^r$, is an expression of the form:

$$t_{n,k} = C_0^n \prod_{i=1}^r C_i^{r_i} \prod_{j=1}^J A_j(n, k)^{\epsilon_j} \tag{8}$$

where C_i are algebraic numbers for $i = 0, \dots, r$, $\epsilon_j = \pm 1$ for $j = 1, \dots, J$, and A_j are integral linear forms in (n, k) that satisfy the *balance condition*:

$$\sum_{j=1}^J \epsilon_j A_j = 0. \tag{9}$$

Remark 1.5. An alternative way of encoding a balanced term is to record the vector (C_0, \dots, C_r) , and the $J \times (r + 2)$ matrix of the coefficients of the linear forms $A_j(n, k)$, and the signs ϵ_j for $j = 1, \dots, r$.

1.5. Balanced terms, generating series and singularities

Given a balanced term t , we will assign a sequence $(a_{t,n})$ and a corresponding generating series $G_t(z)$ to a balanced term t , and will study the set Σ_t of singularities of the analytic continuation of $G_t(z)$ in the complex plane.

Let us introduce some useful notation. Given a linear form $A(n, k)$ in variables (n, k) where $k = (k_1, \dots, k_r)$, and $i = 0, \dots, r$, let us define

$$v_i(A) = a_i, \quad v_0(A) = a_0, \quad \text{where } A(n, k) = a_0n + \sum_{i=1}^r a_i k_i. \tag{10}$$

For $w = (w_1, \dots, w_r)$ we define:

$$A(w) = a_0 + \sum_{i=1}^r a_i w_i. \tag{11}$$

Definition 1.6. Given a balanced term \mathfrak{t} as in (8), define its *Newton polytope* $P_{\mathfrak{t}}$ by

$$P_{\mathfrak{t}} = \{w \in \mathbb{R}^r \mid A_j(w) \geq 0 \text{ for } j = 1, \dots, J\} \subset \mathbb{R}^r. \tag{12}$$

We will assume that $P_{\mathfrak{t}}$ is a compact rational convex polytope in \mathbb{R}^r with non-empty interior. It follows that for every $n \in \mathbb{N}$ we have:

$$\text{support}(\mathfrak{t}_{n,k}) = n P_{\mathfrak{t}} \cap \mathbb{Z}^r. \tag{13}$$

Definition 1.7. Given a balanced term \mathfrak{t} consider the sequence:

$$a_{\mathfrak{t},n} = \sum_{k \in n P_{\mathfrak{t}} \cap \mathbb{Z}^r} \mathfrak{t}_{n,k} \tag{14}$$

(the sum is finite for every $n \in \mathbb{N}$) and the corresponding generating function:

$$G_{\mathfrak{t}}(z) = \sum_{n=0}^{\infty} a_{\mathfrak{t},n} z^n \in \bar{\mathbb{Q}}[[z]]. \tag{15}$$

Here $\bar{\mathbb{Q}}$ denote the field of algebraic numbers. Let $\Sigma_{\mathfrak{t}}$ denote the finite set of singularities of $G_{\mathfrak{t}}$, and $E_{\mathfrak{t}} = \mathbb{Q}(\Sigma_{\mathfrak{t}})$ denote the corresponding number field, following Corollary 1.2.

Remark 1.8. Notice that \mathfrak{t} determines $G_{\mathfrak{t}}$ but *not* vice-versa. Indeed, there are nontrivial identities among multisums of balanced terms. Knowing a complete set of such identities would be very useful in constructing invariants of knotted objects, as well as in understanding relations among periods; see [21].

Remark 1.9. The balance condition of Eq. (9) is imposed so that for every balanced term \mathfrak{t} the corresponding sequence $(a_{\mathfrak{t},n})$ grows at most exponentially. This follows from Stirling’s formula (see Corollary 2.1) and it implies that the power series $G_{\mathfrak{t}}(z)$ is the germ of an analytic function at $z = 0$. Given a proper hypergeometric term $\mathfrak{t}_{n,k}$ in the sense of [33], we can find $\alpha \in \mathbb{Q}$ and $\epsilon = \pm 1$ so that $\mathfrak{t}_{n,k}(\alpha n)!^{\epsilon}$ is a balanced term.

1.6. The definition of $S_{\mathfrak{t}}$ and $K_{\mathfrak{t}}$

Let us observe that if \mathfrak{t} is a balanced term and Δ is a face of its Newton polytope $P_{\mathfrak{t}}$, then $\mathfrak{t}|_{\Delta}$ is also a balanced term.

Definition 1.10. Given a balanced term t as in Eq. (8) consider the following system of *variational equations*:

$$C_i \prod_{j=1}^J A_j(w)^{\epsilon_j v_i(A_j)} = 1 \quad \text{for } i = 1, \dots, r \tag{16}$$

in the variables $w = (w_1, \dots, w_r)$. Let X_t denote the set of complex solutions of (16), with the convention that when $r = 0$ we set $X_t = \{0\}$, and define

$$CV_t = \left\{ C_0^{-1} \prod_{j:A_j(w) \neq 0} A_j(w)^{-\epsilon_j v_0(A_j)} \mid w \in X_t \right\}, \tag{17}$$

$$S_t = \{0\} \cup \cup_{\Delta \text{ face of } P_t} CV_{t|_{\Delta}}, \tag{18}$$

$$K_t = \mathbb{Q}(S_t). \tag{19}$$

Remark 1.11. There are two different incarnations of the set CV_t : it coincides with

- (a) the set of critical values of a potential function; see Theorem 5.
- (b) the evaluation of elements of an additive K -theory group under the entropy regulator function; see Theorem 6.

It is unknown to the author whether X_t is always a finite set. Nevertheless, S_t is always a finite subset of $\bar{\mathbb{Q}}$; see Theorem 5. Eqs. (16) are reminiscent of the *Bethe ansatz*.

1.7. *The conjecture*

Section 1.5 constructs a map:

$$\text{Balanced terms } t \longrightarrow (E_t, \Sigma_t)$$

via generating series $G_t(z)$ and their singularities, where E_t is a number field and Σ_t is a finite subset of E_t . Section 1.6 constructs a map:

$$\text{Balanced terms } t \longrightarrow (K_t, S_t)$$

via solutions of polynomial equations. We are now ready to formulate our main conjecture.

Conjecture 1. *For every balanced term t we have: $\Sigma_t \subset S_t$ and consequently, $E_t \subset K_t$.*

1.8. *Partial results*

Conjecture 1 is known to hold in the following cases:

- (a) For 0-dimensional balanced terms, see Theorem 2.
- (b) For positive special terms, see Theorem 3.
- (c) For 1-dimensional balanced terms, see [13].

Since the finite sets S_t and Σ_t that appear in Conjecture 1 are in principle computable (as explained in Section 1.3), one may try to check random examples. We give some evidence in Section 5. We refer the reader to [16] for an interesting class of 1-dimensional examples related to $6j$ -symbols, and of interest to atomic physics and low-dimensional topology.

The following proposition follows from the classical fact concerning singularities of hypergeometric series; see for example [25, Section 5] and [24].

Theorem 2. *Suppose that $t_{n,k}$ is 0-dimensional balanced term as in (8), with $k = \emptyset$. Then $G_t(z)$ is a hypergeometric series and Conjecture 1 holds.*

When t is positive-dimensional, the generating series $G_t(z)$ is no longer hypergeometric in general. To state our next result, recall that a finite subset $S \subset \mathbb{Q}$ of algebraic numbers is *irreducible* over \mathbb{Q} if the Galois group $\text{Gal}(\mathbb{Q}, \mathbb{Q})$ acts transitively on S .

Definition 1.12. A *special hypergeometric term* $t_{n,k}$ (in short, *special term*) is an expression of the form:

$$t_{n,k} = C_0^n \prod_{i=1}^r C_i^{r_i} \prod_{j=1}^J \binom{B_j(n, k)}{C_j(n, k)} \tag{20}$$

where $C \in \mathbb{Q}, L$ and B_j and C_j are integral linear forms in (n, k) . We will assume that for every $n \in \mathbb{N}$, the support of $t_{n,k} = 0$ as a function of $k \in \mathbb{Z}^r$ is finite. We will call such a term *positive* if $C_i > 0$ for $i = 0, \dots, r$.

Lemma 1.13.

(a) *A balanced term is the ratio of two special terms. In other words, it can always be written in the form:*

$$t_{n,k} = C^{L(n,k)} \prod_{j=1}^s \binom{B_j(n, k)}{C_j(n, k)}^{\epsilon_j} \tag{21}$$

for some integral linear forms B_j, C_j and signs ϵ_j .

(b) *The set of special terms is an abelian monoid with respect to multiplication, whose corresponding abelian group is the set of balanced terms.*

The proof of (a) follows from writing a balanced term in the form:

$$t_{n,k} = C_0^n \prod_{i=1}^r C_i^{r_i} \frac{\prod_{j:\epsilon_j=1}^J A_j(n, k)!}{A(n, k)!} \frac{A(n, k)!}{\prod_{j:\epsilon_j=-1}^J A_j(n, k)!}$$

where

$$A(n, k) = \sum_{j:\epsilon_j=1}^J A_j(n, k) = \sum_{j:\epsilon_j=-1}^J A_j(n, k).$$

To illustrate part (a) of the above lemma for 0-dimensional balanced terms, we have:

$$\frac{(30n)!n!}{(16n)!(10n)!(5n)!} = \binom{30n}{16n} \binom{14n}{10n} \binom{5n}{4n}^{-1}.$$

The above identity also shows that if a balanced term takes integer values, it need not be a special term. This phenomenon was studied by Rodriguez and Villegas; see [29].

Theorem 3. Fix a positive special term $t_{n,k}$ such that $\Sigma_t \setminus \{0\}$ is irreducible over \mathbb{Q} . Then, $\Sigma_t \subset CV_t \subset S_t$ and Conjecture 1 holds.

In a forthcoming publication we will give a proof of Conjecture 1 for 1-dimensional balanced terms; see [13]. Let us end this section with an inverse type (or geometric realization) problem.

Problem 1.14. Given $\lambda \in \bar{\mathbb{Q}}$, does there exist a special term t so that $\lambda \in S_t$?

1.9. Laurent polynomials: A source of special terms

This section, which is of independent interest, associates an special term t_F to a Laurent polynomial F with the property that the generating series $G_{t_F}(z)$ is identified with the trace of the resolvent $R_F(z)$ of F . Combined with Theorem 1, this implies that $R_F(z)$ is a G -function.

If $F \in M_N(\bar{\mathbb{Q}}[x_1^{\pm 1}, \dots, x_d^{\pm 1}])$ is a square matrix of size N with entries Laurent polynomials in d variables, let $\text{Tr}(F)$ denote the constant term of its usual trace. The moment generating series of F is the power series

$$G_F(z) = \sum_{n=0}^{\infty} \text{Tr}(F^n) z^n. \tag{22}$$

Theorem 4.

(a) For every $F \in \bar{\mathbb{Q}}[x_1^{\pm 1}, \dots, x_d^{\pm 1}]$ there exists a special term t_F so that

$$G_F(z) = G_{t_F}(z). \tag{23}$$

Consequently, $G_F(z)$ is a G -function.

- (b) The Newton polytope P_{t_F} depends is a combinatorial simplex which depends only on the monomials that appear in F and not on their coefficients.
- (c) For every $F \in M_N(\bar{\mathbb{Q}}[x_1^{\pm 1}, \dots, x_d^{\pm 1}])$, $G_F(z)$ is a G -function.

We thank C. Sabbah for providing an independent proof of part (c) when $N = 1$ using the regularity of the Gauss–Manin connection. Compare also with [8].

1.10. Plan of the proof

In Section 2, we introduce a potential function associated to a balanced term t and we show that the set of its critical values coincides with the set S_t that features in Conjecture 1. This also implies that S_t is finite.

In Section 3 we assign elements of an extended additive K -theory group to a balanced term \mathfrak{t} , and we show that the set of their values (under the entropy regulator map) coincides with the set $S_{\mathfrak{t}}$ that features in Conjecture 1.

In Section 4 we give a proof of Theorems 3 (using results from hypergeometric functions) and 2 (using an application of Laplace’s method), which are partial case of our Conjecture 1.

In Section 5 we give several examples that illustrate Conjecture 1.

In Section 6 we study a special case of Conjecture 1, with input a Laurent polynomial in many commuting variables.

2. Balanced terms and potential functions

2.1. The Stirling formula and potential functions

As a motivation of a potential function associated to a balanced term, recall *Stirling formula*, which computes the asymptotic expansion of $n!$ (see [25]):

$$\log n! \sim n \log n - n + \frac{1}{2} \log n + \frac{1}{2} \log(2\pi) + O\left(\frac{1}{n}\right). \tag{24}$$

For $x > 0$, we can define $x! = \Gamma(x + 1)$. The Stirling formula implies that for $a > 0$ fixed and $n \in \mathbb{N}$ large, we have:

$$\log(an)! = (n \log n - n)a + a \log(a) + O\left(\frac{\log n}{n}\right). \tag{25}$$

The next corollary motivates our definition of the potential function.

Corollary 2.1. *For every balanced term \mathfrak{t} as in (8) and every w in the interior of $P_{\mathfrak{t}}$ we have:*

$$\mathfrak{t}_{n,nw} = e^{nV_{\mathfrak{t}}(w) + O\left(\frac{\log n}{n}\right)} \tag{26}$$

where the potential function $V_{\mathfrak{t}}$ is defined below.

Definition 2.2. Given a balanced term \mathfrak{t} as in (8), define its corresponding *potential function* $V_{\mathfrak{t}}$ by

$$V_{\mathfrak{t}}(w) = C(w) + \sum_{j=1}^J \epsilon_j A_j(w) \log(A_j(w)) \tag{27}$$

where $w = (w_1, \dots, w_r)$,

$$C(n, k) = \log C_0 \cdot n + \log C_1 \cdot k_1 + \dots + \log C_r \cdot k_r, \tag{28}$$

and

$$C(w) = \log C_0 + \log C_1 w_1 + \dots + \log C_r w_r. \tag{29}$$

$V_{\mathfrak{t}}$ is a multivalued analytic function on the complement of the linear hyperplane arrangement $\mathbb{C}^r \setminus \mathcal{A}_{\mathfrak{t}}$, where

$$\mathcal{A}_{\mathfrak{t}} = \left\{ w \in \mathbb{C}^r \mid \prod_{j=1}^J A_j(w) = 0 \right\}. \tag{30}$$

In fact, let

$$\varpi : \hat{\mathbb{C}}^r \longrightarrow \mathbb{C}^r \setminus \mathcal{A}_{\mathfrak{t}} \tag{31}$$

denote the universal abelian cover of $\mathbb{C}^r \setminus \mathcal{A}_{\mathfrak{t}}$.

The next theorem relates the critical points and critical values of $V_{\mathfrak{t}}$ with the sets $X_{\mathfrak{t}}$ and $CV_{\mathfrak{t}}$ from Definition 1.10.

Theorem 5.

- (a) For every balanced term \mathfrak{t} , $\varpi^{-1}(X_{\mathfrak{t}})$ coincides with the set of critical points of $V_{\mathfrak{t}}$.
- (b) If w is a critical point of $V_{\mathfrak{t}}$, then

$$e^{-V_{\mathfrak{t}}(w)} = C^{-v_0(L)} \prod_{j=1}^J A_j(w)^{-\epsilon_j v_0(A_j)} = C^{-v_0(L)} \prod_{j:A_j(w) \neq 0} A_j(w)^{-\epsilon_j v_0(A_j)}. \tag{32}$$

Thus $CV_{\mathfrak{t}}$ coincides with the exponential of the set of the negatives of the critical values of $V_{\mathfrak{t}}$.

- (c) For every face Δ of the Newton polytope of \mathfrak{t} we have:

$$V_{(\mathfrak{t}|\Delta)} = (V_{\mathfrak{t}})|_{\Delta}.$$

- (d) $S_{\mathfrak{t}}$ is a finite subset of $\bar{\mathbb{Q}}$ and $K_{\mathfrak{t}}$ is a number field.

2.2. Proof of Theorem 5

Let us fix a balanced term \mathfrak{t} as in (8) and a face Δ of its Newton polytope $P_{\mathfrak{t}}$. Without loss of generality, assume that $\Delta = P_{\mathfrak{t}}$. Since

$$\frac{d}{dx} (x \log(x)) = \log(x) + 1 \tag{33}$$

it follows that for every $i = 1, \dots, r$ and every $j = 1, \dots, J$ we have:

$$\frac{\partial}{\partial w_i} A_j(w) \log(A_j(w)) = v_i(A_j) \log(A_j(w)) + v_i(A_j). \tag{34}$$

Adding up with respect to j , using the balancing condition of Eq. (9), and the notation of Section 1.5 applied to the linear form of Eq. (28), it follows that

$$\begin{aligned} \frac{\partial}{\partial w_i} V_t(w) &= \log C_i + \sum_{j=1}^J \epsilon_j v_i(A_j) \log(A_j(w)) + \sum_{j=1}^J \epsilon_j v_i(A_j) \\ &= \log C_i + \sum_{j=1}^J \epsilon_j v_i(A_j) \log(A_j(w)). \end{aligned}$$

This proves that the critical points $w = (w_1, \dots, w_r)$ of V_t are the solutions to the following system of logarithmic variational equations:

$$\log C_i + \sum_{j=1}^J \epsilon_j v_i(A_j) \log(A_j(w)) \in \mathbb{Z}(1) \quad \text{for } i = 1, \dots, r \tag{35}$$

where, for a subgroup K of $(\mathbb{C}, +)$ and an integer $n \in \mathbb{Z}$, we define

$$K(n) = (2\pi i)^n K. \tag{36}$$

Exponentiating, it follows that w satisfies variational equations (16), and concludes the proof of part (a).

For part (b), we will show that if w is a critical point of V_t , the corresponding critical value is given by

$$V_t(w) = \log C_0 + \sum_{j=1}^J \epsilon_j v_0(A_j) \log(A_j(w)) \in \mathbb{C}/\mathbb{Z}(1). \tag{37}$$

Exponentiating, we deduce the first equality of Eq. (32). The second equality follows from the fact that $A_j(w) \neq 0$ for all critical points w of V_t .

To show (37), observe that for any linear form $A(n, k)$ we have:

$$A(w) = v_0(A)w_0 + \sum_{i=1}^r v_i(A)w_i.$$

Suppose that w satisfies logarithmic variational equations (35). Using the definition of the potential function, and collecting terms with respect to w_1, \dots, w_r it follows that

$$\begin{aligned} V_t(w) &= \log C_0 + \sum_{j=1}^J \epsilon_j v_0(A_j) \log(A_j(w)) + \sum_{i=1}^r w_i \left(\log C_i + \sum_{j=1}^J \epsilon_j v_i(A_j) \log(A_j(w)) \right) \\ &= \log C_0 + \sum_{j=1}^J \epsilon_j v_0(A_j) \log(A_j(w)). \end{aligned}$$

This concludes part (b). Part (c) follows from set-theoretic considerations, and part (d) follows from the following facts:

- (i) an analytic function is constant on each component of its set of critical points,

- (ii) the set of critical points are the complex points of an affine variety defined over \mathbb{Q} by (16),
- (iii) every affine variety has finitely many connected components.

This concludes the proof of Theorem 5. \square

3. Balanced terms, the entropy function and additive K -theory

In this section we will assign elements of an extended additive K -theory group to a balanced term, and using them, we will identify our finite set CV_t from Definition 1.10 with the values of the constructed elements under a regulator map; see Theorem 6.

3.1. A brief review of the entropy function and additive K -theory

In this section we will give a brief summary of an extended version of additive K -theory and the entropy function following [15], and motivated by [12]. This section is independent of the rest of the paper, and may be skipped at first reading.

Definition 3.1. Consider the *entropy function* Φ , defined by

$$\Phi(x) = -x \log(x) - (1 - x) \log(1 - x) \tag{38}$$

for $x \in (0, 1)$.

$\Phi(x)$ is a multivalued analytic function on $\mathbb{C} \setminus \{0, 1\}$, given by the double integral of a rational function as follows from

$$\Phi''(x) = -\frac{1}{x} - \frac{1}{1 - x}. \tag{39}$$

For a detailed description of the analytic continuation of Φ , we refer the reader to [15]. Let $\hat{\mathbb{C}}$ denote the *universal abelian cover* of \mathbb{C}^{**} . In [15, Section 1.3] we show that Φ has an analytic continuation:

$$\Phi : \hat{\mathbb{C}} \longrightarrow \mathbb{C}. \tag{40}$$

In [15, Definition 1.7] we show that Φ satisfies three 4-term relations, one of which is the analytic continuation of (48). The other two are dictated by the variation of Φ along the cuts $(1, \infty)$ and $(-\infty, 0)$.

Using the three 4-term relations of Φ , we introduce an extended version $\widehat{\beta}_2(\mathbb{C})$ in [15, Definition 1.7].

Definition 3.2. The *extended group* $\widehat{\beta}_2(\mathbb{C})$ is the \mathbb{C} -vector space generated by the symbols $\langle x \rangle$ with $x = (z; p, q) \in \hat{\mathbb{C}}$, subject to the *extended 4-term relation*:

$$\langle x_0; p_0, q_0 \rangle - \langle x_1; p_1, q_1 \rangle + (1 - x_0) \left\langle \frac{x_1}{1 - x_0}; p_2, q_2 \right\rangle - (1 - x_1) \left\langle \frac{x_0}{1 - x_1}; p_3, q_3 \right\rangle = 0 \tag{41}$$

for $((x_0; p_0, q_0), \dots, (x_3; p_3, q_3)) \in \widehat{4\mathbb{T}}$, and the relations:

$$\langle x; p, q \rangle - \langle x; p, q' \rangle = \langle x; p, q - 2 \rangle - \langle x; p, q' - 2 \rangle, \tag{42}$$

$$\langle x; p, q \rangle - \langle x; p', q \rangle = \langle x; p - 2, q \rangle - \langle x; p' - 2, q \rangle \tag{43}$$

for $x \in \mathbb{C}^{**}$, $p, q, p', q' \in 2\mathbb{Z}$.

Since the three 4-term relations in the definition of $\widehat{\beta_2(\mathbb{C})}$ are satisfied by the entropy function, it follows that Φ gives rise to a regulator map:

$$R: \widehat{\beta_2(\mathbb{C})} \longrightarrow \mathbb{C}. \tag{44}$$

For a motivation of the extended group $\widehat{\beta_2(\mathbb{C})}$ and its relation to additive (i.e., infinitesimal) K -theory and infinitesimal polylogarithms, see [15, Section 1.1] and references therein.

3.2. Balanced terms and additive K -theory

In this section, it will be more convenient to use the presentation (21) of balanced terms. In this case, we have:

$$t_{n,k} = C_0^n \prod_{i=1}^r C_i^{k_i} \prod_{j=1}^J B_j(n, k)!^{\epsilon_j} C_j(n, k)!^{-\epsilon_j} (B_j - C_j)(n, k)!^{-\epsilon_j}. \tag{45}$$

The Stirling formula motivates the constructions in this section. Indeed, we have the following:

Lemma 3.3. For $a > b > 0$, we have:

$$a\Phi\left(\frac{b}{a}\right) = a \log(a) - b \log(b) - (a - b) \log(a - b) \tag{46}$$

and

$$\binom{an}{bn} \sim e^{na\Phi(\frac{b}{a})} \sqrt{\frac{a}{2b(a-b)\pi n}} \left(1 + O\left(\frac{1}{n}\right)\right). \tag{47}$$

Proof. Eq. (46) is elementary. Eq. (47) follows from the Stirling formula (24). \square

The next lemma gives a combinatorial proof of the 4-term relation of the entropy function.

Lemma 3.4. For $a, b, a + b \in (0, 1)$, Φ satisfies the 4-term relation:

$$\Phi(b) - \Phi(a) + (1 - b)\Phi\left(\frac{a}{1 - b}\right) - (1 - a)\Phi\left(\frac{b}{1 - a}\right) = 0. \tag{48}$$

Proof. The 4-term relation follows from the associativity of the multinomial coefficients

$$\binom{\alpha + \beta + \gamma}{\alpha} \binom{\beta + \gamma}{\beta} = \binom{\alpha + \beta + \gamma}{\beta} \binom{\alpha + \gamma}{\alpha} = \frac{(\alpha + \beta + \gamma)!}{\alpha! \beta! \gamma!} \tag{49}$$

applied to $(\alpha, \beta, \gamma) = (an, bn, cn)$ and Lemma 3.3, and the specialization to $a + b + c = 1$. In fact, the 4-term relation (48) and a local integrability assumption uniquely determines Φ up to multiplication by a complex number. See for example, [7] and [1, Section 5.4, p. 66]. \square

Corollary 3.5. *If t is a balanced term as in (21), then its potential function is given by*

$$V_t(w) = C(w) + \sum_{j=1}^J \epsilon_j B_j(w) \Phi \left(\frac{C_j(w)}{B_j(w)} \right). \tag{50}$$

Consider the complement $\mathbb{C}^r \setminus \mathcal{A}'_t$ of the linear hyperplane arrangement given by

$$\mathcal{A}'_t = \left\{ w \in \mathbb{C}^r \mid \prod_{j=1}^J B_j(w) C_j(w) (B_j(w) - 1) (B_j(w) - C_j(w)) = 0 \right\}. \tag{51}$$

Let

$$\varpi : \hat{\mathbb{C}}^r \longrightarrow \mathbb{C}^r \setminus \mathcal{A}'_t \tag{52}$$

denote the universal abelian cover of $\mathbb{C}^r \setminus \mathcal{A}'_t$. For $j = 1, \dots, r$, the functions $B_j(w)$, $C_j(w)/B_j(w)$ have analytic continuation

$$B_j : \hat{\mathbb{C}}^r \longrightarrow \mathbb{C}, \quad \frac{C_j}{B_j} : \hat{\mathbb{C}}^r \longrightarrow \hat{\mathbb{C}}.$$

It follows that the potential function given by (50) has an analytic continuation

$$V_t : \hat{\mathbb{C}}^r \longrightarrow \mathbb{C}. \tag{53}$$

Let \mathcal{C}_t denote the set of critical points of V_t .

Definition 3.6. For a balanced term t as in (21), we define the map

$$\beta_t : \mathcal{C}_t \longrightarrow \widehat{\beta_2(\mathbb{C})} \tag{54}$$

by

$$\beta_t(w) = \sum_{j=1}^J \epsilon_j B_j(w) \left\langle \frac{C_j(w)}{B_j(w)} \right\rangle.$$

Theorem 6. For every balanced term \mathfrak{t} , we have a commutative diagram:

$$\begin{array}{ccc}
 \mathcal{C}_{\mathfrak{t}} & \xrightarrow{\beta_{\mathfrak{t}}} & \widehat{\beta_2(\mathbb{C})} \\
 \downarrow \varpi & & \downarrow e^R \\
 X_{\mathfrak{t}} & \xrightarrow{C_0 e^{-V_{\mathfrak{t}}}} & CV_{\mathfrak{t}}
 \end{array}$$

3.3. Proof of Theorem 6

We begin by observing that the analytic continuation of the logarithm function gives an analytic function

$$\text{Log} : \hat{\mathbb{C}} \longrightarrow \mathbb{C}.$$

The proof of part (a) of Theorem 5 implies that $w \in \mathcal{C}_{\mathfrak{t}}$ if and only if w satisfies the logarithmic variational equations

$$\begin{aligned}
 \log C_i + \sum_{j=1}^J \epsilon_j (v_i(B_j) \text{Log}(B_j(w)) - v_i(C_j) \text{Log}(C_j(w)) \\
 - v_i(B_j - C_j) \text{Log}((B_j - C_j)(w))) = 0
 \end{aligned} \tag{55}$$

for $i = 1, \dots, r$. Exponentiating, this implies that $\mathcal{C}_{\mathfrak{t}} = \varpi^{-1}(X_{\mathfrak{t}})$.

The proof of part (b) of Theorem 5 implies that if $w \in \mathcal{C}_{\mathfrak{t}}$, then

$$\begin{aligned}
 V_{\mathfrak{t}}(w) = \log C_0 + \sum_{j=1}^J \epsilon_j (v_0(B_j) \text{Log}(B_j(w)) - v_0(C_j) \text{Log}(C_j(w)) \\
 - v_0(B_j - C_j) \text{Log}((B_j - C_j)(w))).
 \end{aligned}$$

On the other hand, if $w \in \mathcal{C}_{\mathfrak{t}}$, we have:

$$\begin{aligned}
 R(\beta_{\mathfrak{t}}(w)) &= - \sum_{j=1}^J \epsilon_j B_j(w) \Phi\left(\frac{C_j(w)}{B_j(w)}\right) \\
 &= - \sum_{j=1}^J \epsilon_j (B_j(w) \text{Log}(B_j(w)) - C_j(w) \text{Log}(C_j(w)) \\
 &\quad - (B_j(w) - C_j(w)) \text{Log}((B_j - C_j)(w)))
 \end{aligned}$$

where the last equality follows from the analytic continuation of (46). Expanding the linear forms B_j , C_j , and $B_j - C_j$ with respect to the variables w_i for $i = 0, \dots, r$, and using the logarithmic variational equations (55) (as in the proof of part (b) of Theorem 5), it follows that

$$R(\beta_t(w)) = - \sum_{j=1}^J \epsilon_j (v_0(B_j) \text{Log}(B_j(w)) - v_0(C_j) \text{Log}(C_j(w)) - v_0(B_j - C_j) \text{Log}((B_j - C_j)(w))).$$

Thus,

$$V_t(w) = \log C_0 - R(\beta_t(w)).$$

This concludes the proof of Theorem 6.

4. Proof of Theorems 2 and 3

4.1. Proof of Theorem 2

A 0-dimensional balanced term is of the form

$$t_n = C_0^n \prod_{j=1}^J (b_j n)^{\epsilon_j}$$

where $b_j \in \mathbb{N}$, $\epsilon_j = \pm 1$ for $j = 1, \dots, J$ satisfying $\sum_{j=1}^J \epsilon_j b_j = 0$. Since $\mathbb{Z}^0 = \{0\}$, it follows that

$$a_{t,n} = t_n$$

is so-called closed form. The Newton polytope of t is given by $P_t = \{0\} \subset \mathbb{R}^0$. In addition, $A_j(w) = b_j$ and $v_0(A_j) = b_j$ for $j = 1, \dots, J$. By definition, we have $X_t = \{0\}$, and

$$CV_t = \left\{ C_0^{-1} \prod_{j=1}^J b_j^{-\epsilon_j b_j} \right\}. \tag{56}$$

On the other hand, $G_t(z)$ is a *hypergeometric series* with singularities $\{0, CV_t\}$; see for example [25, Section 5] and [24]. The result follows.

4.2. Proof of Theorem 3

The proof of Theorem 3 is a variant of Laplace’s method and uses the positivity of the restriction of the potential function to P_t . See also [19, Section 5.1.4]. Suppose that $t_{n,k} \geq 0$ for all n, k . Recall the corresponding polytope $P_t \subset \mathbb{R}^f$ and consider the restriction of the potential function to P_t :

$$V_t : P_t \longrightarrow \mathbb{R}.$$

It is easy to show that $\Phi(x) > 0$ for $x \in (0, 1)$. This is illustrated by the plot of the entropy function for $x \in [0, 1]$ (see Fig. 1).




Fig. 1.

It follows that the restriction of the potential function on P_t is nonnegative and continuous. By compactness it follows that the function achieves a maximum in \hat{w} in the interior of P_t . It follows that for every $k \in nP_t \cap \mathbb{Z}^r$ we have:

$$0 \leq t_{n,k} \leq t_{n,n\hat{w}}.$$

Summing up over the lattice points $k \in nP_t \cap \mathbb{Z}^r$, and using the fact that the number of lattice points in a rational convex polyhedron (dilated by n) is a polynomial function of n , it follows that there exist a polynomial $p(n) \in \mathbb{Q}[n]$ so that for all $n \in \mathbb{N}$ we have:

$$t_{n,n\hat{w}} \leq a_{t,n} \leq p(n)t_{n,n\hat{w}}.$$

Using Corollary 2.1, it follows that there exist polynomials $p_1(n), p_2(n) \in \mathbb{Q}[n]$ so that for all $n \in \mathbb{N}$ we have:

$$p_1(n)e^{nV_t(\hat{w})} \leq a_{t,n} \leq p_2(n)e^{nV_t(\hat{w})}.$$

This implies that $G_t(z)$ has a singularity at $z = e^{-V_t(\hat{w})} > 0$. Since the maximum lies in the interior of P_t , it follows that \hat{w} is a critical point of V_t . Thus, $\hat{w} \in S_t$ and consequently, $e^{-V_t(\hat{w})} \in CV_t$.

If in addition $\Sigma_t \setminus \{0\}$ is irreducible over \mathbb{Q} , it follows that the Galois group $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ acts transitively on Σ_t . This implies that $\Sigma_t \subset CV_t$. \square

Remark 4.1. When $t_{n,k} \geq 0$ for all n, k , then $G_t(z)$ has a singularity at $\rho > 0$ where $1/\rho$ is the radius of convergence of $G_t(z)$. This is known as Pringsheim’s theorem, see [32, Section 7.21].

5. Some examples

5.1. A closed form example

As a warm-up example, consider the 1-dimensional special term

$$t_{n,k} = \binom{n}{k} = \frac{n!}{k!(n-k)!}$$

whose corresponding sequence is closed form and the generating series is a rational function:

$$a_{t,n} = 2^n, \quad G_t(z) = \frac{1}{1 - 2z}.$$

In that case $\Sigma_t = \{1/2\}$ and $E_t = \mathbb{Q}$.

To compare with our ansatz, the Newton polytope is given by

$$P_t = [0, 1] \subset \mathbb{R}.$$

Variational equations (16) are

$$\frac{1}{w(1-w)^{-1}} = 1$$

in the variable $w = w_1$, with solution set

$$X_t = \{1/2\}.$$

Thus,

$$CV_t = \left\{ \frac{1}{(1-w)^{-1}} \mid w = 1/2 \right\} = \{1/2\}.$$

For the other two faces $\Delta_0 = \{0\}$ and $\Delta_1 = \{1\}$ of the Newton polytope P_t , the restriction is a 0-dimensional balanced term. Eq. (56) gives that

$$t|_{\Delta_0} = t_{n,k|k=0} = 1, \quad t|_{\Delta_1} = t_{n,k|k=n} = 1.$$

Thus,

$$CV_{t|_{\Delta_0}} = CV_{t|_{\Delta_1}} = \{1\}.$$

This implies that

$$S_t = \{0, 1, 1/2\}, \quad K_t = \mathbb{Q},$$

confirming Conjecture 1. For completeness, the potential function is given by

$$V_t(w) = \Phi(w).$$

5.2. The Apéry sequence

As an illustration of Conjecture 1 and Theorem 3, let us consider the special term

$$t_{n,k} = \binom{n}{k}^2 \binom{n+k}{k}^2 = \left(\frac{(n+k)!}{k!^2(n-k)!} \right)^2 \tag{57}$$

and the corresponding sequence (3). Eq. (5) implies that the singularities of $G(z)$ are a subset of the roots of the equation

$$z^2(z^2 - 34z + 1) = 0.$$

In addition, a nonzero singularity exists. Thus, we obtain that

$$\{0, 17 + 12\sqrt{2}, 17 - 12\sqrt{2}\} \subset \Sigma_t \subset \{0, 17 + 12\sqrt{2}, 17 - 12\sqrt{2}\}, \quad E_t = \mathbb{Q}(\sqrt{2}). \tag{58}$$

Thus E_t is a quadratic number field of type $[2, 0]$ and discriminant 8.

On the other hand, the Newton polytope is given by

$$P_t = [0, 1] \subset \mathbb{R}. \tag{59}$$

Variational equations (16) are

$$\left(\frac{1-w}{w} \frac{1+w}{w}\right)^2 = 1 \tag{60}$$

in the variable $w = w_1$, with solution set

$$X_t = \{1/\sqrt{2}, -1/\sqrt{2}\}.$$

Thus,

$$CV_t = \left\{ \left(\frac{1-w}{1+w}\right)^2 \mid w = \pm 1/\sqrt{2} \right\} = \{17 + 12\sqrt{2}, 17 - 12\sqrt{2}\}.$$

For the other two faces $\Delta_0 = \{0\}$ and $\Delta_1 = \{1\}$ of the Newton polytope P_t the restriction is a 0-dimensional balanced term:

$$t|_{\Delta_0} = t_{n,k} |_{k=0} = 1, \quad t|_{\Delta_1} = t_{n,k} |_{k=n} = \left(\frac{(2n)!}{n!^2}\right)^2.$$

Eq. (56) implies that

$$CV_{t|_{\Delta_0}} = \{1\}, \quad CV_{t|_{\Delta_1}} = \{16\}.$$

Therefore,

$$S_t = \{0, 1, 16, 17 + 12\sqrt{2}, 17 - 12\sqrt{2}\}, \quad K_t = \mathbb{Q}(\sqrt{2}),$$

confirming Conjecture 1. For completeness, the potential function is given by

$$V_t(w) = 2\Phi(w) + 2(1+w)\Phi\left(\frac{w}{1+w}\right).$$

5.3. An example with critical points at the boundary

In this example we find critical points at the boundary of the Newton polytope even though the balanced term is positive. This shows that Theorem 3 is sharp. Consider the balanced term

$$t_{n,k} = \frac{1}{\binom{n}{k}} = \frac{(n-k)!k!}{n!} \tag{61}$$

and the corresponding sequence $(a_{t,n})$. The `zblm` implementation of the WZ algorithm (see [26, 27]) gives that $(a_{t,n})$ satisfies the inhomogeneous recursion relation:

$$-2(n+1)a_{n+1} + (n+2)a_n = -2 - 2n \tag{62}$$

for all $n \in \mathbb{N}$ with initial conditions $a_0 = 1$. It follows that (a_n) satisfies the homogeneous recursion relation:

$$-2(n+3)a_{n+3} + (12+5n)a_{n+2} - 4(n+2)a_{n+1} + (n+2)a_n = 0 \tag{63}$$

for all $n \in \mathbb{N}$ with initial conditions $a_0 = 1, a_1 = 2, a_2 = 5/2$. The corresponding generating series $G_t(z)$ satisfies the inhomogeneous ODE:

$$(z-1)^2(z-2)f'(z) + 2(z-1)^2f(z) + 2 = 0 \tag{64}$$

with initial conditions $f(0) = 1$. We can convert (64) into a homogeneous ODE by differentiating once. It follows that

$$\Sigma_t \subset \{1, 2\}, \quad E_t = \mathbb{Q}. \tag{65}$$

On the other hand, $P_t = [0, 1] \subset \mathbb{R}$. Variational equations (16) are

$$\frac{w}{1-w} = 1$$

in the variable $w = w_1$ with solution set $X_t = \{1/2\}$ and

$$CV_t = \left\{ \frac{1}{1-w} \mid w = \frac{1}{2} \right\} = \{2\}.$$

For the other two faces $\Delta_0 = \{0\}$ and $\Delta_1 = \{1\}$ of the Newton polytope P_t we have:

$$t|_{\Delta_0} = t_{n,k}|_{k=0} = 1, \quad t|_{\Delta_1} = t_{n,k}|_{k=n} = 1.$$

Thus,

$$CV_{t|_{\Delta_0}} = CV_{t|_{\Delta_1}} = \{1\}.$$

This implies that

$$S_t = \{0, 1, 2\}, \quad K_t = \mathbb{Q},$$

confirming Conjecture 1. For completeness, the potential function is given by

$$V_t(w) = -\Phi(w).$$

6. Laurent polynomials and special terms

6.1. Proof of Theorem 4

In this section we will prove Theorem 4. Consider $F \in \bar{\mathbb{Q}}[x_1^{\pm 1}, \dots, x_d^{\pm 1}]$. Let us decompose F into a sum of monomials with coefficients

$$F = \sum_{\alpha \in \mathcal{A}} c_\alpha x^\alpha \tag{66}$$

where \mathcal{A} is the finite set of monomials of F , and where for every $\alpha = (\alpha_1, \dots, \alpha_d)$ we denote $x^\alpha = \prod_{j=1}^d x_j^{\alpha_j}$. Let

$$r = |\mathcal{A}| \tag{67}$$

denote the number of monomials of F . Recall that

$$\Delta_n = \left\{ (y_1, \dots, y_{n+1}) \in \mathbb{R}^{n+1} \mid \sum_{j=1}^{n+1} y_j = 1, y_i \geq 0, i = 1, \dots, n+1 \right\}$$

denotes the standard n -dimensional simplex in \mathbb{R}^{n+1} . Part (a) of Theorem 4 follows easily from the multinomial coefficient theorem. Indeed, for every $n \in \mathbb{N}$ we have:

$$\begin{aligned} F^n &= \sum_{\sum_{\alpha \in \mathcal{A}} k_\alpha = n} \frac{n!}{\prod_{\alpha \in \mathcal{A}} k_\alpha!} \prod_{\alpha \in \mathcal{A}} c_\alpha^{k_\alpha} x^{k_\alpha \alpha} \\ &= \sum_{\sum_{\alpha \in \mathcal{A}} k_\alpha = n} \frac{n!}{\prod_{\alpha \in \mathcal{A}} k_\alpha!} \prod_{\alpha \in \mathcal{A}} c_\alpha^{k_\alpha} \cdot x^{\sum_{\alpha \in \mathcal{A}} k_\alpha \alpha}. \end{aligned}$$

It follows that for every $n \in \mathbb{N}$ we have

$$\begin{aligned} \text{Tr}(F^n) &= \sum_{\sum_{\alpha \in \mathcal{A}} k_\alpha = n, \sum_{\alpha \in \mathcal{A}} k_\alpha \alpha = 0} \frac{n!}{\prod_{\alpha \in \mathcal{A}} k_\alpha!} \prod_{\alpha \in \mathcal{A}} c_\alpha^{k_\alpha} \\ &= \sum_{k \in n P_{t_F} \cap \mathbb{Z}^r} t_{F,n,k} \end{aligned}$$

where

$$t_{F,n,k} = \frac{n!}{\prod_{\alpha \in \mathcal{A}} k_\alpha!} \prod_{\alpha \in \mathcal{A}} c_\alpha^{k_\alpha} \tag{68}$$

and the Newton polygon P_{t_F} is given by

$$P_{t_F} = \Delta_{r-1} \cap W \tag{69}$$

where

$$W = \left\{ (x_\alpha) \in \mathbb{R}^r \mid \sum_{\alpha \in \mathcal{A}} x_\alpha \alpha = 0 \right\} \tag{70}$$

is a linear subspace of \mathbb{R}^r .

To prove part (b) of Theorem 4, let us assume that the origin is in the interior of the Newton polytope of F . Such F are also called *convenient* in singularity theory [22]. If F is not convenient, we can replace it with its restriction F_f to a face f of its Newton polytope that contains the origin and observe that $\text{Tr}((F_f)^n) = \text{Tr}(F^n)$.

When F is convenient, it follows that W has dimension $r - d$ and $W \cap C^o \neq \emptyset$, where C^o is the interior of the cone C which is spanned by the coordinate vectors in \mathbb{R}^r . (b) follows from Lemma 6.3 below.

For part (c) of Theorem 4, fix $F \in M_N(\bar{\mathbb{Q}}[x_1^{\pm 1}, \dots, x_d^{\pm 1}])$, $G_F(z)$. Recall that for an invertible matrix A we have $A^{-1} = \det(A)^{-1} \text{Cof}(A)$, where $\text{Cof}(A)$ is the co-factor matrix. It follows that

$$\sum_{n=0}^{\infty} F^n z^n = (I - zF)^{-1} = \frac{1}{\det(I - zF)} \text{Ad}(I - zF) \tag{71}$$

where $\text{Ad}(I - zF) \in M_N(\bar{\mathbb{Q}}[x_1^{\pm 1}, \dots, x_d^{\pm 1}, z])$. Now

$$\det(I - zF) = 1 - \sum_{\alpha \in \mathcal{S}} c_\alpha z^{d_\alpha} x^\alpha$$

where \mathcal{S} is a finite set, $c_\alpha \in \bar{\mathbb{Q}}$ and $d_\alpha \in \mathbb{N} \setminus \{0\}$. Thus,

$$\frac{1}{\det(I - zF)} = \sum_{n=0}^{\infty} \sum_{\sum_{\alpha \in \mathcal{S}} k_\alpha = n} \frac{n!}{\prod_{\alpha \in \mathcal{S}} k_\alpha!} z^{\sum_{\alpha \in \mathcal{S}} d_\alpha k_\alpha} \prod_{\alpha \in \mathcal{S}} x^{\sum_{\alpha \in \mathcal{S}} k_\alpha \alpha}.$$

Substituting the above into Eq. (71) and taking the constant term, it follows that there exists a finite set S and a finite collection $t_F^{(j)}$ of balanced terms for $j \in S$ such that for every $n \in \mathbb{N}$ we have:

$$\text{Tr}(F^n) = \sum_{j \in S} a_{t_F^{(j)}, n}.$$

It follows that the moment generating series $G_F(z)$ (defined in Eq. (22)) is given by

$$G_F(z) = \sum_{j \in S} G_{t_F^{(j)}}(z).$$

Since $G_{t_F^{(j)}}(z)$ is a G -function (by Theorem 1), and the set of G -functions is closed under addition (see [2]), this concludes the proof Theorem 4.

Remark 6.1. The balanced terms $t_F^{(j)}$ in the above proof use affine linear forms, rather than linear ones. In other words, they are given by

$$t_{n,k} = C_0^n \prod_{i=1}^r C_i^{r_i} \prod_{j=1}^J A_j(n, k)^{\epsilon_j} \tag{72}$$

where C_i are algebraic numbers for $i = 0, \dots, r$, $\epsilon_j = \pm 1$ for $j = 1, \dots, J$, and A_j are affine linear forms, given by $A_j(n, k) = A_j^{\text{lin}}(n, k) + b_j$ where $A_j^{\text{lin}}(n, k)$ are linear forms that satisfy the balance condition

$$\sum_{j=1}^J \epsilon_j A_j^{\text{lin}} = 0. \tag{73}$$

Theorem 1 remains true for such balanced terms. For an balanced term t of the form (72) consider the balanced term t^{lin} defined by

$$t_{n,k}^{\text{lin}} = C_0^n \prod_{i=1}^r C_i^{r_i} \prod_{j=1}^J A_j^{\text{lin}}(n, k)^{\epsilon_j}. \tag{74}$$

We define $S^t = S^{t^{\text{lin}}}$.

Remark 6.2. In the notation of Theorem 4, P_{t_F} is a *simple* polytope and the associated toric variety is projective and has *quotient singularities*; see [11]. In other words, the stabilizers of the torus action on the toric variety are finite abelian groups.

The following lemma was communicated to us by J. Pommersheim.

Lemma 6.3. *If W is a linear subspace of \mathbb{R}^n of dimension s , and $W \cap C^o \neq \emptyset$, where C is the cone spanned by the n coordinate vectors in \mathbb{R}^n , then $\Delta_{n-1} \cap W$ is a combinatorial simplex in V .*

Proof. We can prove the claim by downward induction on s . When $s = n - 1$, W is a hyperplane. Write $D_{n-1} = C \cap H$ where H is the hyperplane given by $\sum_{i=1}^n x_i = 0$. Observe that C is a simplicial cone and the intersection $C \cap W$ is a simplicial cone in W . Since the intersection of a simplicial cone inside C with H is a simplicial cone, it follows that $D_{n-1} \cap W = (C \cap W) \cap H$ is a simplicial cone. This proves the claim when $s = n - 1$. A downward induction on s concludes the proof. \square

6.2. The Newton polytope of a Laurent polynomial and its associated balanced term

In a later publication we will study the close relationship between the Newton polytope of a Laurent polynomial F and the Newton polytope of the corresponding special term t_F . In what follows, fix a Laurent polynomial F as in (66) and its associated balanced term t_F as in (68). With the help of the commutative diagram of Proposition 6.5 below, we will compare the extended critical values of F with the set S_{t_F} . Keep in mind that [8] use the Newton polytope of F , whereas our ansatz uses the Newton polytope of t_F .

Consider the polynomial map

$$\phi : \bar{\mathbb{Q}}[w_\alpha^{\pm 1} \mid \alpha \in \mathcal{A}] \longrightarrow \bar{\mathbb{Q}}[x_1^{\pm 1}, \dots, x_d^{\pm 1}] \tag{75}$$

given by $\phi(w_\alpha) = x^\alpha$. Its kernel $\ker(\phi)$ is a *monomial ideal*. Adjoin the monomial w_{α_0} to \mathcal{A} (if it not already there), where $\alpha_0 = 0 \in \mathbb{Z}^d$, and consider the homogeneous ideal $\ker^h(\phi)$, where the degree of w_α is α .

Lemma 6.4. *Variational equations (16) for the balanced term t_F are equivalent to the following system of equations:*

$$\begin{cases} \prod_{\alpha \in \mathcal{A}} x_\alpha^{-p_\alpha} c_\alpha^{p_\alpha} = 1 & \text{for } (w_\alpha^{p_\alpha}) \in \ker^h(\phi), \\ \sum_{\alpha \in \mathcal{A}} x_\alpha \alpha = 0, \\ \sum_{\alpha \in \mathcal{A}} x_\alpha = 1 \end{cases} \tag{76}$$

in the variables $(x_\alpha)_{\alpha \in \mathcal{A}}$.

Proof. This follows easily from the description of the Newton polytope P_{t_F} and the shape of the balanced term t . \square

Consider the rational function

$$\Psi_F : (\mathbb{C}^*)^d \setminus F^{-1}(0) \longrightarrow (\mathbb{C}^*)^r, \quad u \mapsto \Psi_F(u) = \left(\frac{c_\alpha u^\alpha}{F(u)} \right)_{\alpha \in \mathcal{A}}. \tag{77}$$

Observe that the image of Ψ_F lies in the complex affine simplex

$$\left\{ (x_\alpha) \in (\mathbb{C}^*)^r \mid \sum_{\alpha \in \mathcal{A}} x_\alpha = 1 \right\}.$$

Proposition 6.5.

- (a) *If $u = (u_1, \dots, u_d)$ is a critical point of the restriction of F on $(\mathbb{C}^*)^d$ with nonvanishing critical value, then $\Psi_F(u)$ satisfies variational equations (16) for the maximal face P_{t_F} of the Newton polytope P_{t_F} of t_F .*
- (b) *Restricting to those critical points, we have a commutative diagram:*

$$\begin{array}{ccc} \text{Critical points of } F \text{ on } (\mathbb{C}^*)^d & \xrightarrow{\Psi_F} & X_{t_F, P_{t_F}} \\ \downarrow F & & \downarrow \text{Map (17)} \\ \text{Critical values of } F & \longrightarrow & S_{t_F, P_{t_F}} \end{array} \tag{78}$$

- (c) *The top horizontal map of the above diagram is 1–1 and onto.*

Proof. For (a) we will use the alternative system of variational equations given in (76). Suppose that $u = (u_1, \dots, u_d)$ is a critical point of F with nonzero critical value, and let $(w_\alpha)_{\alpha \in \mathcal{A}}$ denote the tuple $(c_\alpha u^\alpha / F(u))_{\alpha \in \mathcal{A}}$. We need to show that $(w_\alpha)_{\alpha \in \mathcal{A}}$ is a solution to Eq. (76). Consider a point $(p_\alpha)_{\alpha \in \mathcal{A}} \in P_{t_F}$. Then, we can verify the first line of Eq. (76) as follows

$$\begin{aligned} \prod_{\alpha \in \mathcal{A}} w_\alpha^{-p_\alpha} c_\alpha^{p_\alpha} &= \prod_{\alpha \in \mathcal{A}} \left(\frac{c_\alpha u^\alpha}{F(u)} \right)^{-p_\alpha} c_\alpha^{p_\alpha} \\ &= u^{-\sum_{\alpha \in \mathcal{A}} p_\alpha \alpha} F(u)^{-\sum_{\alpha \in \mathcal{A}} p_\alpha} \\ &= 1 \end{aligned}$$

since $(w_\alpha^{p_\alpha})_{\alpha \in \mathcal{A}}$ in the kernel of ϕ . To verify the second line of Eq. (76), recall that

$$z \partial_z (z^k) = k z^k.$$

Since u is a critical point of F , it follows that for every $i = 1, \dots, d$ we have

$$u_i \partial_{u_i} F(u) = \sum_{\alpha \in \mathcal{A}} v_i(\alpha) c_\alpha u^\alpha = 0$$

where $v_i(\alpha)$ is the i th coordinate of α . It follows that $\sum_{\alpha \in \mathcal{A}} w_\alpha \alpha = 0$, which verifies the second line of (76). To verify the third line, we compute:

$$\sum_{\alpha \in \mathcal{A}} \frac{c_\alpha u^\alpha}{F(u)} = \frac{1}{F(u)} \sum_{\alpha \in \mathcal{A}} c_\alpha u^\alpha = \frac{F(u)}{F(u)} = 1.$$

(b) and (c) follow from similar computations. \square

Let us end this section with an example that illustrates Lemma 6.4 and Proposition 6.5.

Example 6.6. Consider the Laurent polynomial

$$F(x_1, x_2) = ax_1 + bx_1^{-1} + cx_2 + dx_2^{-1} + f x_1 x_2 + g.$$

Its Newton polytope is given in Fig. 2.

With the notation from the previous section, we have $d = 2, r = 6$. Moreover, for every $n \in \mathbb{N}$, we have

$$F^n = \sum_{k_1 + \dots + k_6 = n} \frac{n!}{k_1! \dots k_6!} a^{k_1} b^{k_2} c^{k_3} d^{k_4} f^{k_5} g^{k_6} x_1^{k_1 - k_2 + k_5} x_2^{k_3 - k_4 + k_5}.$$

It follows that

$$\text{Tr}(F^n) = \sum_{(k_1, \dots, k_6) \in n P_{t_F} \cap \mathbb{Z}^6} \frac{n!}{k_1! \dots k_6!} a^{k_1} b^{k_2} c^{k_3} d^{k_4} f^{k_5} g^{k_6}$$

where the Newton polytope $P_{t_F} \subset \mathbb{R}^6$ is given by




Fig. 2.

$$P_{t_F} = \{(x_1, \dots, x_6) \in \mathbb{R}^6 \mid x_1 - x_2 + x_5 = 0, x_3 - x_4 + x_5 = 0, x_1 + \dots + x_6 = 1, x_i \geq 0\}. \tag{79}$$

We can use the variables (x_1, x_3, x_5) to parametrize P_{t_F} as follows:

$$P_{t_F} = \{(x_1, x_3, x_5) \in \mathbb{R}^3 \mid 1 \geq 2x_1 + 2x_3 + 3x_5, x_i \geq 0\}. \tag{80}$$

This confirms that P_{t_F} is a combinatorial 2-dimensional simplex.

If $(k_1, \dots, k_6) \in P_{t_F}$, then $(k_2, k_4, k_6) = (k_1 + k_5, k_3 + k_5, n - 2k_1 - 2k_3 - 3k_5)$, and

$$\begin{aligned} \text{Tr}(F^n) &= \sum_{2k_1+2k_3+2k_5=n} \frac{n!}{k_1!(k_1+k_5)!k_3!(k_3+k_5)!k_5!(n-2k_1-2k_3-3k_5)!} \\ &\quad \times \left(\frac{ab}{g^2}\right)^{k_1} \left(\frac{cd}{g^2}\right)^{k_3} \left(\frac{bdf}{g^3}\right)^{k_5} g^n. \end{aligned}$$

Variational equations (16) are:

$$\begin{aligned} \frac{1}{w_1(w_1+w_5)(1-2w_1-2w_3-3w_5)^{-2}} \frac{ab}{g^2} &= 1, \\ \frac{1}{w_3(w_3+w_5)(1-2w_1-2w_3-3w_5)^{-2}} \frac{cd}{g^2} &= 1, \\ \frac{1}{(w_1+w_5)(w_3+w_5)w_5(1-2w_1-2w_3-2w_5)^{-3}} \frac{bdf}{g^3} &= 1, \end{aligned} \tag{81}$$

in the variables (w_1, w_3, w_5) . Reintroducing the variables w_2, w_4 and w_6 defined by

$$w_2 = w_1 + w_5, \quad w_4 = w_3 + w_5, \quad w_6 = 1 - 2w_1 - 2w_3 - 2w_5,$$

variational equations become

$$\begin{aligned} w_2 &= w_1 + w_5, \\ w_4 &= w_3 + w_5, \\ w_6 &= 1 - 2w_1 - 2w_3 - 2w_5, \end{aligned} \tag{82}$$

$$\begin{aligned}\frac{1}{w_1 w_2 w_6^{-2}} \frac{ab}{g^2} &= 1, \\ \frac{1}{w_3 w_4 w_6^{-2}} \frac{cd}{g^2} &= 1, \\ \frac{1}{w_2 w_4 w_5 w_6^{-3}} \frac{bdf}{g^3} &= 1.\end{aligned}\tag{83}$$

On the other hand, $w_{\alpha_0} = w_6$ and $\ker(\phi)$ is generated by the relations $w_6 = 1$, $w_1 w_2 = 1$, $w_3 w_4 = 1$ and $w_2 w_4 w_5 = 1$ which lead to the homogeneous relations $w_1 w_2 w_6^{-2} = 1$, $w_3 w_4 w_6^{-2} = 1$ and $w_2 w_4 w_5 w_6^{-3}$ for $\ker^h(\phi)$ that appear in the above variational equations. This illustrates Lemma 6.4.

Suppose now that $u = (u_1, u_2)$ is a critical point of F . Then, u satisfies the equations:

$$\frac{\partial F}{\partial u_1} = a - \frac{b}{u_1^2} + f u_2 = 0,\tag{84}$$

$$\frac{\partial F}{\partial u_2} = c - \frac{d}{u_2^2} + f u_1 = 0.\tag{85}$$

The map Ψ_F is defined by $\Psi_F(u_1, u_2) = (w_1, \dots, w_6)$ where

$$\begin{aligned}w_1 &= \frac{a u_1}{F(u)}, & w_2 &= \frac{b}{u_1 F(u)}, & w_3 &= \frac{c u_2}{F(u)}, \\ w_4 &= \frac{d}{u_2 F(u)}, & w_5 &= \frac{f u_1 u_2}{F(u)}, & w_6 &= \frac{g}{F(u)}.\end{aligned}$$

If $u = (u_1, u_2)$ satisfies Eqs. (84), (85), it is easy to see that $w = \Psi_F(u)$ satisfies variational equations (82) and (83). For example, we have

$$w_3 + w_5 - w_4 = \frac{c u_2}{F(u)} + \frac{f u_1 u_2}{F(u)} - \frac{d}{u_2 F(u)} = \frac{u_2}{F(u)} \left(c + f u_1 - \frac{d}{u_2^2} \right) = 0$$

and

$$\frac{1}{w_3 w_4 w_6^{-2}} \frac{cd}{g^2} = \frac{1}{\frac{c u_2}{F(u)} \frac{d}{u_2 F(u)} \left(\frac{g}{F(u)} \right)^{-2}} \frac{cd}{g^2} = 1.$$

This illustrates Proposition 6.5.

Acknowledgments

The author wishes to thank Y. André, N. Katz, M. Kontsevich, J. Pommersheim, C. Sabbah, and especially D. Zeilberger for many enlightening conversations, R.I. van der Veen for a careful reading of the manuscript, and the anonymous referee for comments that improved the exposition. An early version of the paper was presented in an Experimental Mathematics Seminar in Rutgers in the spring of 2007. The author wishes to thank D. Zeilberger for his hospitality.

References

- [1] J. Aczél, J. Dhombres, *Functional Equations in Several Variables*, Encyclopedia Math. Appl., vol. 31, Cambridge Univ. Press, Cambridge, 1989.
- [2] Y. André, *G-functions and Geometry*, Aspects Math., vol. E13, Vieweg, Braunschweig, 1989.
- [3] Y. André, Séries Gevrey de type arithmétique. I. Théorèmes de pureté et de dualité, *Ann. of Math. (2)* 151 (2000) 705–740.
- [4] G. Birkhoff, W. Trjitzinsky, Analytic theory of singular difference equations, *Acta Math.* 60 (1932) 1–89.
- [5] D.V. Chudnovsky, G.V. Chudnovsky, Applications of Padé approximations to the Grothendieck conjecture on linear differential equations, in: *Number Theory*, in: *Lecture Notes in Math.*, vol. 1135, Springer-Verlag, 1985, pp. 52–100.
- [6] O. Costin, S. Garoufalidis, Resurgence of the Kontsevich–Zagier power series, *Ann. Inst. Fourier (Grenoble)*, in press.
- [7] Z. Daróczy, Generalized information functions, *Inform. Control* 16 (1970) 36–51.
- [8] J.J. Duistermaat, W. van der Kallen, Constant terms in powers of a Laurent polynomial, *Indag. Math.* 9 (1998) 221–231.
- [9] B. Dwork, G. Gerotto, F.J. Sullivan, *An Introduction to G-Functions*, Ann. Math. Stud., vol. 133, Princeton Univ. Press, Princeton, NJ, 1994.
- [10] P. Flajolet, R. Sedgewick, *Analytic combinatorics: Functional equations, rational and algebraic functions*, electronic book.
- [11] W. Fulton, *Introduction to Toric Varieties*, Ann. Math. Stud., vol. 131, Princeton Univ. Press, Princeton, NJ, 1993.
- [12] S. Garoufalidis, q -terms, singularities and the extended Bloch group, preprint, arXiv: 0708.0018, 2007.
- [13] S. Garoufalidis, Resurgence of 1-dimensional hypergeometric multisums, preprint, 2007.
- [14] S. Garoufalidis, G -functions and multisum versus holonomic sequences, preprint, arXiv: 0708.4354, 2007.
- [15] S. Garoufalidis, An extended version of additive K-theory, J. K-theory, in press.
- [16] S. Garoufalidis, R. van der Veen, Asymptotics of $6j$ -symbols, preprint, 2008.
- [17] R. Jungen, Sur les séries de Taylor n'ayant que des singularités algébriques-logarithmiques sur leur cercle de convergence, *Comment. Math. Helv.* 3 (1931) 266–306.
- [18] N.M. Katz, Nilpotent connections and the monodromy theorem: Applications of a result of Turrittin, *Inst. Hautes Études Sci. Publ. Math.* 39 (1970) 175–232.
- [19] D. Knuth, *The Art of Computer Programming*, vol. 3, Sorting and Searching, Addison–Wesley, Reading, MA, 1973.
- [20] M. Kontsevich, Yu. Manin, Gromov–Witten classes, quantum cohomology, and enumerative geometry, in: *Mirror Symmetry, II*, in: *AMS/IP Stud. Adv. Math.*, vol. 1, Amer. Math. Soc., Providence, RI, 1999, pp. 607–653.
- [21] M. Kontsevich, Yu. Manin, D. Zagier, Periods, in: *Mathematics Unlimited—2001 and Beyond*, 2001, pp. 771–808.
- [22] A.G. Kouchirenko, Polyèdres de Newton et nombres de Milnor, *Invent. Math.* 32 (1976) 1–31.
- [23] N. Nilsson, Some growth and ramification properties of certain integrals on algebraic manifolds, *Ark. Mat.* 5 (1965) 463–476.
- [24] N.E. Norlund, Hypergeometric functions, *Acta Math.* 94 (1955) 289–349.
- [25] F. Olver, *Asymptotics and Special Functions*, reprint A K Peters Classics, AK Peters, Wellesley, MA, 1997.
- [26] P. Paule, A. Riese, A Mathematica q -analogue of Zeilberger's algorithm based on an algebraically motivated approach to q -hypergeometric telescoping, in: *Special Functions, q -Series and Related Topics*, in: *Fields Inst. Commun.*, vol. 14, Amer. Math. Soc., Providence, RI, 1997, pp. 179–210.
- [27] P. Paule, A. Riese, Mathematica software: <http://www.risc.uni-linz.ac.at/research/combinat/risc/software/qZeil/>.
- [28] M. Petkovšek, H.S. Wilf, D. Zeilberger, *A = B*, AK Peters, Wellesley, MA, 1996.
- [29] F. Rodriguez-Villegas, Integral ratios of factorials and algebraic hypergeometric functions, talk in Oberwolfach, math.NT/0701362.
- [30] C.L. Siegel, Über einige Anwendungen Diophantischer Approximationen, *Abh. Preuss. Akad. Wiss.* 1 (1929) 1–70; reprinted in: *Gesam. Abh.* 1 (16) (1966) 209–266.
- [31] R. Stanley, *Enumerative Combinatorics*, vols. 1, 2, second ed., Cambridge Univ. Press, Cambridge, 1997.
- [32] E.C. Titchmarsh, *The Theory of Functions*, second ed., Oxford Univ. Press, Oxford, 1939.
- [33] H. Wilf, D. Zeilberger, An algorithmic proof theory for hypergeometric (ordinary and q) multisum/integral identities, *Invent. Math.* 108 (1992) 575–633.
- [34] J. Wimp, D. Zeilberger, Resurrecting the asymptotics of linear recurrences, *J. Math. Anal. Appl.* 111 (1985) 162–176.
- [35] D. Zeilberger, A holonomic systems approach to special functions identities, *J. Comput. Appl. Math.* 32 (1990) 321–368.