


Periodic Approximations for Energy Spectra in Aperiodic Media

Jean BELLISSARD

*Westfälische Wilhelms-Universität, Münster
Department of Mathematics*

*Georgia Institute of Technology, Atlanta
School of Mathematics & School of Physics
e-mail: jeanbel@math.gatech.edu*

Sponsoring



*SFB 878, Münster,
Germany*

Contributors

G. DE NITTIS, *Departamento de Matemática, Pontificia Universidad Católica de Chile, Santiago, Chile*

S. BECKUS, *Department of Mathematics, Technion, Israel*

H. D. CORNEAN, *Institut for Matematiske Fag, Aalborg Universitet, Aalborg, Denmark*

S. BECKUS,
Spectral approximation of aperiodic Schrödinger operators,
PhD Thesis, U. Jena, October 6, 2016
arXiv: 1610.05894

Recent Work

S. BECKUS, J. BELLISSARD,
Continuity of the Spectrum of a Field of Self-Adjoint Operators,
Ann. Henri Poincaré, **17**, (2016), 3425-3442.

S. BECKUS, J. BELLISSARD, G. DE NITTIS,
Spectral Continuity for Aperiodic Quantum Systems I. General Theory,
arXiv: 1709.00975,

S. BECKUS, J. BELLISSARD, G. DE NITTIS,
Spectral Continuity for Aperiodic Quantum Systems II. Periodic approximations in 1D,
in preparation

S. BECKUS, J. BELLISSARD, H. CORNEAN,
Spectral Lipschitz Continuity for FLC Aperiodic Systems,
in preparation

S. BECKUS, J. BELLISSARD,
Spectral Stability of Schrödinger operators for the Kohmoto model,
in preparation

S. BECKUS, J. BELLISSARD, G. DE NITTIS,
Computing the Spectrum of 1D-tight binding Models,
in preparation

Main References

J. E. ANDERSON, I. PUTNAM,
Topological invariants for substitution tilings and their associated C^ -algebras,*
Ergodic Theory Dynam. Systems, **18**, (1998), 509-537.

F. GÄHLER, Talk given at *Aperiodic Order, Dynamical Systems, Operator Algebra and Topology*
Victoria, BC, August 4-8, 2002, *unpublished*.

J. BELLISSARD, R. BENEDETTI, J. M. GAMBAUDO,
Spaces of Tilings, Finite Telescopic Approximations,
Comm. Math. Phys., **261**, (2006), 1-41.


J. BELLISSARD, *Wannier Transform for Aperiodic Solids*, Talks given at
EPFL, Lausanne, June 3rd, 2010
KIAS, Seoul, Korea September 27, 2010
Georgia Tech, March 16th, 2011
Cergy-Pontoise September 5-6, 2011
U.C. Irvine, May 15-19, 2013
WCOAS, UC Davis, October 26, 2013
Simons Center for Geometry and Physics, U. Stonybrook, June 7, 2016
Dep. Math, U. Carlos III, Madrid, November 2, 2016
ICMS, Edinburgh, June 26-30, 2017
online at <http://people.math.gatech.edu/~jeanbel/talksjbE.html>

Content

1. Motivation
2. Continuous Fields
3. Conclusion

I - Motivations

Motivation



Physica Scripta. Vol. T9, 193–198, 1985

Renormalization of Quasiperiodic Mappings

Stellan Ostlund and Seung-hwan Kim

Spectrum of the Kohmoto model

(Fibonacci Hamiltonian)

$$(H\psi)(n) = \psi(n+1) + \psi(n-1) + \lambda \chi_{(0,\alpha]}(x - n\alpha) \psi(n)$$

as a function of α .

Method:
transfer matrix calculation

Motivation




Fig. 3. – We show, respectively, the IDOS of the Octonacci chain (up) and the IDOS of the labyrinth, for a) $r = 0.8$ (no gap, finite measure), b) $r = 0.6$ (some gaps and finite measure) and c) $r = 0.3$ (infinity of gaps and zero measure). The energy varies between -2 and 2 , since $r < 1$.

C. SIRE


**Electronic Spectrum of a 2D Quasi-Crystal Related
to the Octagonal Quasi-Periodic Tiling.**

EUROPHYSICS LETTERS

Europhys. Lett., 10 (5), pp. 483-488 (1989)

Solvable 2D-model, reducible to 1D-calculations

Motivation



A sample of the icosahedral quasicrystal AlPdMn

Methodologies

- For *one dimensional Schrödinger* equation of the form

$$H\psi(x) = -\frac{d^2\psi}{dx^2} + V(x)\psi(x)$$

a *transfer matrix* approach has been used for a long time to analyze the spectral properties (*Bogoliubov '36*).

- A *KAM-type* perturbation theory has been used successfully (*Dinaburg, Sinai '76, JB '80's*).

Methodologies

- For *discrete* one-dimensional models of the form

$$H\psi(n) = t_{n+1}\psi(n+1) + t_n\psi(n-1) + V_n\psi(n)$$

a *transfer matrix approach* is the most efficient method, both for numerical calculation and for mathematical approach:

- the *KAM-type* perturbation theory also applies (*JB '80's*).
- models defined by substitutions using the *trace map*
(*Khomoto et al., Ostlund et al. '83, JB '89, JB, Bovier, Ghez, Damanik... in the nineties*)
- theory of cocycle (*Avila, Jitomirskaya, Damanik, Krikorian, Gorodetsky...*).

Methodologies

- In higher dimension almost no rigorous results were available
- Exceptions are for models that are Cartesian products of 1D models (*Sire '89, Damanik, Gorodestky, Solomyak '14*)
- Numerical calculations performed on quasi-crystals have shown that
 - Finite cluster calculation lead to a large number of *spurious edge states*.
 - *Periodic approximations* are much more efficient
 - Some periodic approximations exhibit *defects* giving *contributions* in the energy spectrum. (*JB, Iochum, Scoppola, Testard '89*)

II - Continuous Fields

Continuous Fields of Hamiltonians

$A = (A_t)_{t \in T}$ is a *field of self-adjoint operators* whenever

1. T is a topological space,
2. for each $t \in T$, \mathcal{H}_t is a Hilbert space,
3. for each $t \in T$, A_t is a self-adjoint operator acting on \mathcal{H}_t .

The field $A = (A_t)_{t \in T}$ is called *p^2 -continuous* whenever, for every polynomial $p \in \mathbb{R}(X)$ with degree at most 2, the following norm map is *continuous*

$$\Phi_p : t \in T \mapsto \|p(A_t)\| \in [0, +\infty)$$

Continuous Fields of Hamiltonians

Theorem: *(S. Beckus, J. Bellissard '16)*

- 1. A field $A = (A_t)_{t \in T}$ of self-adjoint bounded operators is p_2 -continuous if and only if the spectrum of A_t , seen as a compact subset of \mathbb{R} , is a continuous function of t with respect to the Hausdorff metric.*
- 2. Equivalently $A = (A_t)_{t \in T}$ is p_2 -continuous if and only if the spectral gap edges of A_t are continuous functions of t .*

Continuous Fields of Hamiltonians

The field $A = (A_t)_{t \in T}$ is called *p_2 - α -Hölder continuous* whenever, for every polynomial $p \in \mathbb{R}(X)$ with degree at most 2, the following norm map is *α -Hölder continuous*

$$\Phi_p : t \in T \mapsto \|p(A_t)\| \in [0, +\infty)$$


uniformly w.r.t. $p(X) = p_0 + p_1X + p_2X^2 \in \mathbb{R}(X)$ such that $|p_0| + |p_1| + |p_2| \leq M$, for some $M > 0$.

Continuous Fields of Hamiltonians

Theorem: *(S. Beckus, J. Bellissard '16)*


- 1. A field $A = (A_t)_{t \in T}$ of self-adjoint bounded operators is p_2 - α -Hölder continuous then the spectrum of A_t , seen as a compact subset of \mathbb{R} , is an $\alpha/2$ -Hölder continuous function of t with respect to the Hausdorff metric.*
- 2. In such a case, the edges of a spectral gap of A_t are α -Hölder continuous functions of t at each point t where the gap is open.*
- 3. At any point t_0 for which a spectral gap of A_t is closing, if the tip of the gap is isolated from other gaps, then its edges are $\alpha/2$ -Hölder continuous functions of t at t_0 .*
- 4. Conversely if the gap edges are α -Hölder continuous, then the field A is p_2 - α -Hölder continuous.*

Continuous Fields of Hamiltonians



*The spectrum of the Harper model
the Hamiltonian is p -Lipshitz continuous*

(JB, '94)



A gap closing (enlargement)

Continuous Fields on C^* -algebras

(Tomiyama 1958, Dixmier-Douady 1962)

Given a topological space T , let $\mathcal{A} = (\mathcal{A}_t)_{t \in T}$ be a family of C^* -algebras.

A *vector field* is a family $a = (a_t)_{t \in T}$ with $a_t \in \mathcal{A}_t$ for all $t \in T$.

\mathcal{A} is called *continuous* whenever there is a family Υ of vector fields such that,

- for all $t \in T$, the set Υ_t of elements a_t with $a \in \Upsilon$ is a *dense *-subalgebra* of \mathcal{A}_t
- for all $a \in \Upsilon$ the map $t \in T \mapsto \|a_t\| \in [0, +\infty)$ is *continuous*
- a vector field $b = (b_t)_{t \in T}$ belongs to Υ if and only if, for any $t_0 \in T$ and any $\epsilon > 0$, there is U an open neighborhood of t_0 and $a \in \Upsilon$, with $\|a_t - b_t\| < \epsilon$ whenever $t \in U$.

Continuous Fields on C^* -algebras

Theorem *If \mathcal{A} is a continuous field of C^* -algebras and if $a \in \mathcal{Y}$ is a continuous self-adjoint vector field, then, for any continuous function $f \in C_0(\mathbb{R})$, the maps $t \in T \mapsto \|f(a_t)\| \in [0, +\infty)$ are continuous*

In particular, such a vector field is p_2 -continuous

Groupoids

(Ramsay '76, Connes, 79, Renault '80)

A *groupoid* G is a category the object of which G_0 and the morphism of which G make up two sets. More precisely

- there are two maps $r, s : G \rightarrow G_0$ (*range* and *source*)
- $(\gamma, \gamma') \in G_2$ are *compatible* whenever $s(\gamma) = r(\gamma')$
- there is an associative *composition law* $(\gamma, \gamma') \in G_2 \mapsto \gamma \circ \gamma' \in G$, such that $r(\gamma \circ \gamma') = r(\gamma)$ and $s(\gamma \circ \gamma') = s(\gamma')$
- a *unit* e is an element of G such that $e \circ \gamma = \gamma$ and $\gamma' \circ e = \gamma'$ whenever compatibility holds; then $r(e) = s(e)$ and the map $e \rightarrow x = r(e) = s(e) \in G_0$ is a *bijection* between units and objects;
- each $\gamma \in G$ admits an *inverse* such that $\gamma \circ \gamma^{-1} = r(\gamma) = s(\gamma^{-1})$ and $\gamma^{-1} \circ \gamma = s(\gamma) = r(\gamma^{-1})$

Locally Compact Groupoids

- A groupoid G is *locally compact* whenever
 - G is endowed with a locally compact Hausdorff topology
 - the maps r, s , the *composition* and the *inverse* are *continuous* functions.

Then the set of units is a closed subset of G .

- A *Haar system* is a family $\lambda = (\lambda^x)_{x \in G_0}$ of positive Borel measures on the fibers $G^x = r^{-1}(x)$, such that
 - if $\gamma : x \rightarrow y$, then $\gamma^* \lambda^x = \lambda^y$
 - if $f \in C_c(G)$ is continuous with compact support, then the map $x \in G_0 \mapsto \lambda^x(f)$ is *continuous*.

Locally Compact Groupoids

Locally Compact Groupoids

Example:

Let Ω be a *compact Hausdorff* space, let G be a *locally compact group* acting on Ω by homeomorphisms. Then $\Gamma = \Omega \times G$ becomes a locally compact groupoid as follows

- $\Gamma_0 = \Omega$, is the set of *units*,
- $r(\omega, g) = \omega$ and $s(\omega, g) = g^{-1}\omega$
- $(\omega, g) \circ (g^{-1}\omega, h) = (\omega, gh)$
- Each fiber $\Gamma^\omega \simeq G$, so that if μ is the *Haar measure* on G , it gives a Haar system λ with $\lambda^\omega = \mu$ for all $\omega \in \Omega$.

This groupoid is called the *crossed-product* and is denoted $\Omega \rtimes G$

Groupoid C^* -algebra

Let G be a locally compact groupoid with a Haar system λ . Then the complex vector space space $C_c(G)$ of complex valued continuous functions with compact support on G becomes a $*$ -algebra as follows

- **Product (convolution):**

$$ab(\gamma) = \int_{G^x} a(\gamma') b(\gamma'^{-1} \circ \gamma) d\lambda^x(\gamma') \quad x = r(\gamma)$$

- **Adjoint:**

$$a^*(\gamma) = \overline{a(\gamma^{-1})}$$

Groupoid C^* -algebra

The following construction gives a C^* -norm

- for each $x \in G_0$, let $\mathcal{H}_x = L^2(G^x, \lambda^x)$
- for $a \in C_c(G)$, let $\pi_x(a)$ be the operator on \mathcal{H}_x defined by

$$\pi_x(a)\psi(\gamma) = \int_{G^x} a(\gamma^{-1} \circ \gamma') \psi(\gamma') d\lambda^x(\gamma')$$

- $(\pi_x)_{x \in G_0}$ gives a faithful covariant family of $*$ -representations of $C_c(G)$, namely if $\gamma : x \rightarrow y$ then $\pi_x \sim \pi_y$.
- then $\|a\| = \sup_{x \in G_0} \|\pi_x(a)\|$ is a C^* -norm; the completion of $C_c(G)$ with respect to this norm is called the *reduced C^* -algebra* of G and is denoted by $C_{red}^*(G)$.

Continuous Fields of Groupoids

(N. P. Landsman, B. Ramazan, 2001)

- A *field of groupoid* is a triple (G, T, p) , where G is a groupoid, T a set and $p : G \rightarrow T$ a map, such that, if $p_0 = p \upharpoonright_{G_0}$, then $p = p_0 \circ r = p_0 \circ s$
- Then the subset $G_t = p^{-1}\{t\}$ is a groupoid depending on t .
- If G is *locally compact*, T a *Hausdorff* topological space and p *continuous* and *open*, then $(G, T, P) = (G_t)_{t \in T}$ is called a *continuous field of groupoids*.

Continuous Fields of Groupoids

Theorem: *(N. P. Landsman, B. Ramazan, 2001)*

Let (G, T, p) be a continuous field of locally compact groupoids with Haar systems. If G_t is amenable for all $t \in T$, then the field $\mathcal{A} = (\mathcal{A}_t)_{t \in T}$ of C^ -algebras defined by $\mathcal{A}_t = C^*(G_t)$ is continuous.*

III - Tautological Groupoid

Periodic Approximations

Approximating an aperiodic system by a periodic one makes sense within the following framework

- Ω is a *compact* Hausdorff metrizable space,
- a locally compact *group* G acts on Ω by homeomorphisms,
- $\mathcal{J}(\Omega)$ is the set of *closed G -invariant* subsets of Ω :
 - a subset $M \in \mathcal{J}(\Omega)$ is *minimal* if all its G -orbits are dense.
 - a point $\omega \in \Omega$ is called *periodic* if there is a *uniform lattice* $\Lambda \subset G$ such that $g\omega = \omega$ for $g \in \Lambda$. In such a case $\text{Orb}(\omega)$ is a quotient of G/Λ , and is thus compact.
 - if G is *discrete*, any periodic orbit is a *finite set*.

Periodic Approximations

Example 1)- Subshifts

Let \mathcal{A} be a finite set (alphabet). Let $\Omega = \mathcal{A}^{\mathbb{Z}}$: it is compact for the product topology. The shift operator S defines a \mathbb{Z} -action.

1. A sequence $\xi = (x_n)_{n \in \mathbb{Z}}$ is *periodic* if and only if ξ can be written as an infinite repetition of a finite word. The S -orbit of ξ is then *finite*.
2. The set of periodic points of Ω is *dense*.
3. A *subshift* is provided by a closed S -invariant subset, namely a point in $\mathcal{J}(\Omega)$.

Periodic Approximations

Example 2)- Delone Sets A *Delone* set $\mathcal{L} \subset \mathbb{R}^d$ is

- a discrete closed subset,
- there is $0 < r$ such that each ball of radius r intersects \mathcal{L} at one point *at most*,
- here is $0 < R$ such that each ball of radius R intersects \mathcal{L} at one point *at least*.

Then

1. the set $\Omega = \text{Del}_{r,R}$ of such Delone sets in \mathbb{R}^d can be endowed with a topology that makes it *compact*,
2. the group \mathbb{R}^d *acts* on Ω by homeomorphisms,
3. the periodic Delone sets make up a *dense subset* in Ω .

Periodic Approximations

Question: *in which sense can one approximate a minimal infinite G -invariant subset by a sequence of periodic orbits ?*

The Fell and Vietoris Topologies

(Vietoris 1922, Fell 1962)

Given a *topological space* X , let $\mathcal{C}(X)$ be the set of *closed subsets* of X .

Let $F \subset X$ be closed and let \mathcal{F} be a finite family of open sets. Then

$$\mathcal{U}(F, \mathcal{F}) = \{G \in \mathcal{C}(X); G \cap F = \emptyset, G \cap O \neq \emptyset, \forall O \in \mathcal{F}\}$$

Then the family of $\mathcal{U}(F, \mathcal{F})$'s is a basis for a topology called the *Vietoris topology*.

Replacing F by a compact set K , the same definition leads to the *Fell topology*.

The Fell and Vietoris Topologies

- $\mathcal{C}(X)$ is *Fell-compact*,
- if X is locally compact and Hausdorff, $\mathcal{C}(X)$ is *Hausdorff* for both Fell and Vietoris,
- if (X, d) is a *complete metric* space, the Vietoris topology coincides with the topology defined by the *Hausdorff metric*.
- If X is *compact* both topologies *coincide*.

Theorem *If (Ω, G) is a topological dynamical system, the set $\mathcal{J}(\Omega)$ is compact for both the Fell and the Vietoris topologies.*

The Fell and Vietoris Topologies

Example:

If $(\Omega = \mathcal{A}^{\mathbb{Z}}, S)$, periodic orbits *ARE NOT* Vietoris-dense in $\mathcal{J}(\Omega)$

For instance, if $\mathcal{A} = \{0, 1\}$ let $\xi_0 \in \Omega$ be the sequence defined by

$$\xi_0 = (x_n)_{n \in \mathbb{Z}} \quad x_n = \begin{cases} 0 & \text{if } n < 0 \\ 1 & \text{if } n \geq 0 \end{cases}$$

Then $\text{Orb}(\xi)$ is *isolated* in $\mathcal{J}(\Omega)$ for the Vietoris topology.

The Tautological Groupoid

- Let $\mathcal{T}(\Omega) \subset \mathcal{J}(\Omega) \times \Omega$ be the set of pairs (M, ω) such that $\omega \in M$. Endowed with the product topology it is *compact Hausdorff*.
- G *acts* on it by homeomorphisms through $g(M, \omega) = (M, g\omega)$.
- Let $\Gamma = \mathcal{T}(\Omega) \rtimes G$, let $T = \mathcal{J}(\Omega)$ and let $p : \Gamma \rightarrow T$ defined by

$$p(M, \omega, g) = M$$

Then (G, T, p) is a *continuous field* of locally compact groupoids.

- If G is *amenable*, then the family $\mathcal{A} = (\mathcal{A}_M)_{M \in \mathcal{J}(\Omega)}$ where $\mathcal{A}_M = C^*(\Gamma_M)$ gives a continuous field of C^* -algebras.

The Tautological Groupoid

Hence

If M is a closed invariant subset of Ω that is a Vietoris-limit point of the set of periodic orbit, then any continuous field of Hamiltonian in \mathcal{A} has a spectrum that can be approximated by the spectrum of a suitable sequence of periodic approximations.

Question: *Which invariant subsets of Ω are Vietoris limit points of periodic orbits ?*

IV - Periodic Approximations for Subshifts

Subshifts


Let \mathcal{A} be a finite *alphabet*, let $\Omega = \mathcal{A}^{\mathbb{Z}}$ be equipped with the shift S . Let $\Sigma \in \mathcal{J}(\Omega)$ be a subshift. Then

- given $l, r \in \mathbb{N}$ an *(l, r) -collared dot* is a dotted word of the form $u \cdot v$ with u, v being words of length $|u| = l, |v| = r$ such that uv is a *sub-word* of at least one element of Σ
- an *(l, r) -collared letter* is a dotted word of the form $u \cdot a \cdot v$ with $a \in \mathcal{A}, u, v$ being words of length $|u| = l, |v| = r$ such that uav is a sub-word of at least one element of Σ : *a collared letter links two collared dots*
- let $\mathcal{V}_{l,r}$ be the set of (l, r) -collared dots, let $\mathcal{E}_{l,r}$ be the set of (l, r) -collared letters: then the pair $\mathcal{G}_{l,r} = (\mathcal{V}_{l,r}, \mathcal{E}_{l,r})$ gives a finite *directed graph*


(de Bruijn, '46, Anderson-Putnam '98, Gähler, '01)

The Fibonacci Tiling

- **Alphabet:** $\mathcal{A} = \{a, b\}$
- **Fibonacci sequence:** generated by the *substitution* $a \rightarrow ab, b \rightarrow a$ starting from either $a \cdot a$ or $b \cdot a$




Left: $\mathcal{G}_{1,1}$



Right: $\mathcal{G}_{8,8}$

The Thue-Morse Tiling


- **Alphabet:** $\mathcal{A} = \{a, b\}$
- **Thue-Morse sequences:** generated by the *substitution* $a \rightarrow ab, b \rightarrow ba$ starting from either $a \cdot a$ or $b \cdot a$



Thue-Morse $\mathcal{G}_{1,1}$

The Rudin-Shapiro Tiling


- **Alphabet:** $A = \{a, b, c, d\}$
- **Rudin-Shapiro sequences:** generated by the *substitution* $a \rightarrow ab, b \rightarrow ac, c \rightarrow db, d \rightarrow dc$ starting from either $b \cdot a, c \cdot a$ or $b \cdot d, c \cdot d$




Rudin-Shapiro $\mathcal{G}_{1,1}$

The Full Shift on Two Letters

- **Alphabet:** $\mathcal{A} = \{a, b\}$ all possible word allowed.



$\mathcal{G}_{1,2}$



$\mathcal{G}_{2,2}$

Strongly Connected Graphs

The de Bruijn graphs are

- *simple*: between two vertices there is at most one edge,
- *connected*: if the sub-shift is *topologically transitive*, (i.e. one orbit is dense), then between any two vertices, there is at least one path connected them,
- has *no dangling vertex*: each vertex admits at least one ingoing and one outgoing vertex,
- if $n = l + r = l' + r'$ then the graphs $\mathcal{G}_{l,r}$ and $\mathcal{G}_{l',r'}$ are *isomorphic* and denoted by \mathcal{G}_n .

Strongly Connected Graphs

(S. Beckus, PhD Thesis, 2016)

A directed graph is called *strongly connected* if any pair x, y of vertices there is an *oriented path* from x to y and another one from y to x .

Proposition: *If the sub-shift Σ is minimal (i.e. every orbit is dense), then each of the de Bruijn graph is strongly connected.*

Main result:

Theorem: *A subshift $\Sigma \subset A^{\mathbb{Z}}$ can be Vietoris approximated by a sequence of periodic orbits if and only if it admits a sequence of strongly connected de Bruijn graphs.*

Open Problem

Question:

Is there a similar criterion for the space of Delone sets in \mathbb{R}^d or for some remarkable subclasses of it ?

Some *sufficient conditions* have been found for $\Omega = \mathcal{A}^G$, where G is a discrete, countable and *amenable group*, in particular when $G = \mathbb{Z}^d$.

(S. Beckus, PhD Thesis, 2016)



Thanks for listening !