

# ON UNIFORM CONVERGENCE OF DIAGONAL MULTIPOINT PADÉ APPROXIMANTS FOR ENTIRE FUNCTIONS

D. S. LUBINSKY

ABSTRACT. We prove that for most entire functions  $f$  in the sense of category, a strong form of the Baker-Gammel-Wills Conjecture holds. More precisely, there is an infinite sequence  $\mathcal{S}$  of positive integers  $n$ , such that given any  $r > 0$ , and multipoint Padé approximants  $R_n$  to  $f$  with interpolation points in  $\{z : |z| \leq r\}$ ,  $\{R_n\}_{n \in \mathcal{S}}$  converges locally uniformly to  $f$  in the plane. The sequence  $\mathcal{S}$  does not depend on  $r$ , nor on the interpolation points. For entire functions with smooth rapidly decreasing coefficients, full diagonal sequences of multipoint Padé approximants converge.

Padé approximation, Multipoint Padé approximants, spurious poles.  
41A21, 41A20, 30E10.

## 1. INTRODUCTION<sup>1</sup>

Let  $D$  be an open connected subset of  $\mathbb{C}$ , and  $f : D \rightarrow \mathbb{C}$  be analytic. Given  $n \geq 0$  and  $2n + 1$  not necessarily distinct points  $\Lambda_n = \{z_j\}_{j=1}^{2n+1}$  in  $D$ , and

$$\omega_n(z) = \omega_n(\Lambda_n, z) = \prod_{j=1}^{2n+1} (z - z_j),$$

the  $(n, n)$  multipoint Padé approximant to  $f$  with interpolation set  $\Lambda_n$  is a rational function

$$R_n(\Lambda_n, z) = \frac{p_n(\Lambda_n, z)}{q_n(\Lambda_n, z)},$$

or more simply,

$$R_n(z) = \frac{p_n(z)}{q_n(z)},$$

---

*Date:* August 1, 2017.

<sup>1</sup>Research supported by NSF grant DMS1362208

where  $p_n$  and  $q_n$  are polynomials of degree  $\leq n$ , with  $q_n$  not identically zero, such that

$$\frac{f(z)q_n(z) - p_n(z)}{\omega_n(z)}$$

is analytic in  $D$ . The special case where all  $z_j = 0$ , gives the Padé approximant  $[n/n](z)$ . It is easily seen that  $R_n$  exists and is unique, though  $p_n$  and  $q_n$  are not separately unique.

The convergence of Padé and multipoint Padé approximants is a much studied subject, with uniform convergence established for large classes of special functions. One of the pitfalls of the method is the appearance of **spurious poles**, namely poles that do not reflect the analytic properties of the interpolated function  $f$  [1], [2], [5], [8], [12], [14], [22], [23], [24], [27], [29], [30], [31], [33]. For this reason, the most general results, such as the Nuttall-Pommerenke theorem, often involve convergence in capacity, rather than uniform convergence. In 1961, Baker, Gammel, and Wills nevertheless conjectured that at least a subsequence of the diagonal Padé sequence converges locally uniformly. Throughout this paper,

$$B_r = \{z : |z| < r\}, r > 0.$$

### Baker-Gammel-Wills Conjecture

*Let  $f$  be meromorphic in  $B_1$  and analytic at 0. Then there is a subsequence  $\{[n/n]\}_{n \in \mathcal{S}}$  of  $\{[n/n]\}_{n \geq 1}$  that converges uniformly to  $f$  in compact subsets of  $B_1$  omitting poles of  $f$ .*

The author showed in 2001 [24] that the conjecture is false, by considering the Rogers-Ramanujan function with a non-standard value of  $q$  on the unit circle. While this provided a meromorphic counterexample, A.P. Buslaev quickly followed [6] with an analytic counterexample, formed from an algebraic function, and then showed that even the Rogers-Ramanujan function provides an analytic counterexample [7]. George Baker [3] subsequently noted that for these counterexamples, just two subsequences together provide locally uniform convergence in the unit ball. He went on to conjecture that a patchwork of finitely many subsequences can provide locally uniform convergence for functions meromorphic in the ball [4].

One of the unsolved issues is whether the Baker-Gammel-Wills conjecture is valid for entire functions, or perhaps even functions meromorphic in the plane. To date, there is still no counterexample. The author proved [19] that the Baker-Gammel-Wills conjecture is true for most entire functions in the sense of category.

In this paper, we shall show that a stronger form of the conjecture, allowing interpolation points in any compact set, with the same subsequence, is true for most entire functions in the sense of category. Accordingly, let  $\mathcal{A}$  denote the space of entire functions, with metric defined in terms of power series coefficients: if

$$f(z) = \sum_{j=0}^{\infty} a_j z^j \text{ and } g(z) = \sum_{j=0}^{\infty} b_j z^j,$$

then define

$$(1.1) \quad d^*(f, g) = \sup_{j \geq 0} |a_j - b_j|^{1/\max\{j, 1\}}.$$

Convergence in this space is equivalent to uniform convergence in compact sets. Recall that a subset of  $\mathcal{A}$  is of the first category, if it is a countable union of nowhere dense sets. As such, it is small in the sense of category. Recall too that an  $F_\sigma$  set is a countable union of closed sets.

**Theorem 1.1**

*There is an  $F_\sigma$  subset  $\mathcal{E}$  of  $\mathcal{A}$  of the first category, such that for  $f \in \mathcal{A} \setminus \mathcal{E}$ , there is an infinite subsequence  $\mathcal{S}$  of positive integers with the following property: given any  $r > 0$  and for  $n \in \mathcal{S}$ , multipoint Padé approximants  $R_n$  to  $f$  of type  $(n, n)$  formed from interpolation points  $\Lambda_n \subset B_r$ , we have*

$$(1.2) \quad \lim_{n \rightarrow \infty, n \in \mathcal{S}} R_n(z) = f(z)$$

*uniformly in compact subsets of the plane.*

Observe that while  $\mathcal{S}$  depends on  $f$ , it does not depend on the ball  $B_r$  in which the interpolation points lie. As far as the author is aware,  $e^z$  is the only function for which diagonal rational interpolants with interpolation points in any compact set (and that are not restricted to include complex conjugate interpolation points) had been proven to converge locally uniformly [10], [32]. For Markov-Stieltjes functions, convergence of diagonal multipoint Padé approximants, with interpolation points symmetric about the real axis, has been investigated in [9], [13].

We also prove some more explicit results when the Maclaurin series coefficients decay rapidly and/or smoothly:

**Theorem 1.2**

Assume that

$$(1.3) \quad f(z) = \sum_{j=0}^{\infty} a_j z^j$$

where  $a_j \neq 0$  for  $j \geq 0$  and for some fixed  $J$ , and for  $j \geq J$ ,

$$(1.4) \quad \left| \frac{a_{j-1} a_{j+1}}{a_j^2} \right| \leq \chi^2$$

where  $\chi < \rho_0$  and  $\rho_0 = 0.4559\dots$  is the positive root of the equation

$$(1.5) \quad \sum_{j=1}^{\infty} \rho^{j^2} = \frac{1}{2}.$$

Let  $r > 0$ . For  $n \geq 1$ , let  $R_n$  denote a multipoint Padé approximant to  $f$  formed with interpolation points in  $B_r$ . Then uniformly in compact subsets of the plane,

$$\lim_{n \rightarrow \infty} R_n(z) = f(z).$$

**Theorem 1.3**

Assume that  $f$  is given by (1.3), where  $a_j \neq 0$  for  $j \geq 0$  and for some  $|q| < 1$ ,

$$(1.6) \quad \lim_{j \rightarrow \infty} \frac{a_{j-1} a_{j+1}}{a_j^2} = q$$

Then the conclusion of Theorem 1.2 remains valid.

We note that Theorems 1.2 and 1.3 were proved for the special case of Padé approximants in [20], [21]. In [20], the slightly more general condition  $\chi \leq \rho_0$  was allowed. We note also that Theorems 1.2 and 1.3 and the results of [25] show that given  $s > r > 0$ , then for large enough  $n$ ,  $f - R_n$  formed from interpolation points in  $B_r$ , has exactly  $2n+1$  zeros, counting multiplicity, in  $B_s$ . Related results dealing with smooth Maclaurin series coefficients appear in [11], [15], [16], [28]. Without smoothness but with more rapid decay, we prove convergence of a subsequence:

**Theorem 1.4**

Assume that for  $f$  given by (1.3),

$$(1.7) \quad \limsup_{j \rightarrow \infty} |a_j|^{1/j^2} < \frac{1}{3}.$$

Then there is a subsequence  $\mathcal{S}$  of integers with the property (1.2) as described in Theorem 1.1.

Theorem 1.1 suggests a stronger form of the Baker-Gammel-Wills Conjecture for entire functions:

**Conjecture 1.5**

Let  $f$  be entire. Then there is an infinite subsequence  $\mathcal{S}$  of positive integers with the following property: given any  $r > 0$  and for  $n \in \mathcal{S}$ , multipoint Padé approximants  $R_n$  to  $f$  of type  $(n, n)$  formed from interpolation points  $\Lambda_n \subset B_r$ , we have (1.2).

We close this section with more notation, firstly, finite differences: given distinct  $z_1, z_2, z_3, \dots$ , define  $f[z_1] = f(z_1)$ ,

$$f[z_1, z_2] = \frac{f(z_2) - f(z_1)}{z_2 - z_1};$$

and recursively, for  $r \geq 2$ ,

$$f[z_1, \dots, z_{r+1}] = \frac{f[z_1, \dots, z_{r-1}, z_{r+1}] - f[z_1, \dots, z_{r-1}, z_r]}{z_{r+1} - z_r}.$$

When points coalesce, that is not all  $\{z_j\}$  are distinct, the finite difference is defined as the appropriate derivative. We also set

$$(1.8) \quad f_{i,j} = f[z_{i+1}, z_{i+2}, \dots, z_{j+1}].$$

We shall make extensive use of the following formula for the denominator in  $R_n(\Lambda_n, z)$  when  $\Lambda_n = \{z_j\}_{j=1}^{2n+1}$  [2, p. 339]:

$$\det \begin{bmatrix} f_{n,n+1} & f_{n,n+2} & \cdots & f_{n,2n} & \prod_{k=1}^n (z - z_k) \\ f_{n-1,n+1} & f_{n-1,n+2} & \cdots & f_{n-1,2n} & \prod_{k=1}^{n-1} (z - z_k) \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ f_{0,n+1} & f_{0,n+2} & \cdots & f_{0,2n} & 1 \end{bmatrix}.$$

It is valid as long as this last determinant is not identically 0. By row and column swaps, we can recast it (absorbing a sign change into the

numerator polynomial) as

(1.9)

$$q_n(\Lambda_n, z) = \det \begin{bmatrix} 1 & f_{0,n+1} & f_{0,n+2} & \cdots & f_{0,2n} \\ z - z_1 & f_{1,n+1} & f_{1,n+2} & \cdots & f_{1,2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \prod_{k=1}^{n-1} (z - z_k) & f_{n-1,n+1} & f_{n-1,n+2} & \cdots & f_{n-1,2n} \\ \prod_{k=1}^n (z - z_k) & f_{n,n+1} & f_{n,n+2} & \cdots & f_{n,2n} \end{bmatrix}.$$

Throughout this paper, we assume that  $f$  is entire, not a polynomial, and has Maclaurin series given by (1.3). The paper is organized as follows: we prove Theorems 1.1 and 1.4 in Section 2, Theorem 1.2 in Section 3, and Theorem 1.3 in Section 4.

## 2. PROOF OF THEOREMS 1.1 AND 1.4

We begin by bounding coefficients of non-polynomial entire functions, much as in [19]. Let

$$K = \max \{1, |a_0|\}.$$

Define an increasing sequence of integers

$$0 = j_0 < j_1 < j_2 < \dots$$

and positive numbers  $\{\rho_j\}$  as follows: first, choose  $j_1 \geq 1$  such that

$$\rho_{j_1}^{-1} = \left( \frac{|a_{j_1}|}{K} \right)^{1/j_1} = \max \left\{ \left( \frac{|a_j|}{K} \right)^{1/j} : j \geq 1 \right\}.$$

Having defined  $\rho_{j_1}, \dots, \rho_{j_k}$ , define  $\rho_{j_{k+1}}$  by

$$\rho_{j_{k+1}}^{-1} = \left| \frac{a_{j_{k+1}}}{a_{j_k}} \right|^{\frac{1}{j_{k+1} - j_k}} = \max \left\{ \left| \frac{a_j}{a_{j_k}} \right|^{\frac{1}{j - j_k}} : j > j_k \right\}.$$

If there is more than one choice of  $j_k$ , choose the largest one. Define

$$\rho_n = \rho_{j_{k+1}} \text{ for } j_k + 1 \leq n \leq j_{k+1} \text{ and } k \geq 0.$$

### Lemma 2.1

(a)

$$(2.1) \quad |a_n| \leq K / \prod_{\ell=1}^n \rho_\ell, \text{ for } n \geq 0,$$

with equality when

$$n = j_k \text{ for some } k \geq 1.$$

- (b)  $\{\rho_k\}$  is an increasing sequence with limit  $\infty$ .  
(c) If  $n = j_k$  for some  $k \geq 1$ , then for  $r \geq -n$ ,

$$(2.2) \quad \left| \frac{a_{n+r}}{a_n} \right| \leq \begin{cases} \rho_{n+1}^{-r}, & r \geq 0 \\ \rho_n^{-r}, & r < 0 \end{cases}.$$

**Proof**

(a) Suppose that  $k$  is given, and  $j_k + 1 \leq n \leq j_{k+1}$ . Then by definition of  $j_{k+1}$ ,

$$\begin{aligned} \left| \frac{a_n}{a_{j_k}} \right|^{\frac{1}{n-j_k}} &\leq \rho_{j_{k+1}}^{-1} \\ \Rightarrow |a_n| &\leq |a_{j_k}| \rho_{j_{k+1}}^{-(n-j_k)} = |a_{j_k}| / \prod_{\ell=j_k+1}^n \rho_\ell. \end{aligned}$$

We have equality if  $n = j_{k+1}$ . Applying this inequality recursively to  $a_{j_k}, a_{j_{k-1}}, \dots$ , we obtain (2.1), with equality if  $n$  equals some  $j_k$ , some  $k \geq 1$ .

(b) Now as  $j_{k+2} > j_{k+1}$ ,

$$\begin{aligned} \rho_{j_{k+1}}^{-1} &= \left| \frac{a_{j_{k+1}}}{a_{j_k}} \right|^{\frac{1}{j_{k+1}-j_k}} > \left| \frac{a_{j_{k+2}}}{a_{j_k}} \right|^{\frac{1}{j_{k+2}-j_k}} \\ &= \left| \frac{a_{j_{k+2}}}{a_{j_{k+1}}} \right|^{\frac{1}{j_{k+2}-j_k}} \left| \frac{a_{j_{k+1}}}{a_{j_k}} \right|^{\frac{1}{j_{k+2}-j_k}} \\ \Rightarrow \left| \frac{a_{j_{k+1}}}{a_{j_k}} \right|^{\frac{1}{j_{k+1}-j_k} - \frac{1}{j_{k+2}-j_k}} &> \left| \frac{a_{j_{k+2}}}{a_{j_{k+1}}} \right|^{\frac{1}{j_{k+2}-j_k}} \\ \Rightarrow \left| \frac{a_{j_{k+1}}}{a_{j_k}} \right|^{\frac{j_{k+2}-j_{k+1}}{j_{k+1}-j_k}} &> \left| \frac{a_{j_{k+2}}}{a_{j_{k+1}}} \right| \\ \Rightarrow \left| \frac{a_{j_{k+1}}}{a_{j_k}} \right|^{\frac{1}{j_{k+1}-j_k}} &> \left| \frac{a_{j_{k+2}}}{a_{j_{k+1}}} \right|^{\frac{1}{j_{k+2}-j_{k+1}}} \\ \Rightarrow \rho_{j_{k+1}}^{-1} &> \rho_{j_{k+2}}^{-1}. \end{aligned}$$

The monotonicity of  $\{\rho_n\}$  follows, and the fact that  $f$  is entire, forces them to have limit  $\infty$ .

(c) If  $r \geq 0$ , we have (using that there is equality in (2.1) for  $n = j_k$ ),

$$\left| \frac{a_{n+r}}{a_n} \right| \leq \prod_{\ell=n+1}^{n+r} \rho_\ell^{-1} \leq \rho_{n+1}^{-r}.$$

If  $r < 0$ , we instead have

$$\left| \frac{a_{n+r}}{a_n} \right| \leq \prod_{\ell=n+r}^{n-1} \rho_\ell \leq \rho_n^{-r}.$$

■

We shall frequently use a series expansion for finite differences. Assume  $R > 0$ ,  $\ell \geq 1$ , and  $z_1, z_2, \dots, z_\ell \in \overline{B_R}$ . Then by the contour integral representation for finite differences [26, p. 11], if  $\Gamma = \{t : |t| = S\}$ , where  $S > R$ , then

$$\begin{aligned} f[z_1, z_2, \dots, z_\ell] &= \frac{1}{2\pi i} \int_\Gamma \frac{f(t)}{\prod_{k=1}^{\ell} (t - z_k)} dt \\ &= \frac{1}{2\pi i} \int_\Gamma \frac{f(t)}{t^\ell} \sum_{i_1 \dots i_\ell \geq 0} \frac{z_1^{i_1} z_2^{i_2} \dots z_\ell^{i_\ell}}{t^{i_1+i_2+\dots+i_\ell}} dt \\ &= \sum_{i_1 \dots i_\ell \geq 0} z_1^{i_1} z_2^{i_2} \dots z_\ell^{i_\ell} a_{\ell-1+i_1+i_2+\dots+i_\ell}. \end{aligned}$$

(2.3)

**Lemma 2.2**

Assume that  $n = j_i$  for some  $i \geq 1$ , where the  $\{j_i\}$  are as above. Assume that all  $|z_j| \leq R$  and  $\rho_n > R$ .

(a) Then for  $0 \leq j, k \leq n$ , with the notation (1.8),

$$(2.4) \quad (\rho_n \rho_{n+1})^{\frac{k-j}{2}} \left| \frac{f_{j,n+k}}{a_n} \right| \leq \left( \frac{\rho_n}{\rho_{n+1}} \right)^{\frac{|k-j|}{2}} (1 - R/\rho_n)^{-(n+1+k-j)}.$$

(b) For  $0 \leq j \leq n$ ,

$$(2.5) \quad \left| \frac{f_{j,n+j}}{a_n} - 1 \right| \leq (1 - R/\rho_{n+1})^{-(n+1)} - 1.$$



**Proof**

(a) Let  $\ell = n + 1 + k - j$ . From the series (2.3) above

$$\begin{aligned}
 \left| \frac{f_{j,n+k}}{a_n} \right| &= \left| \sum_{i_1 \dots i_\ell \geq 0} z_{j+1}^{i_1} z_{j+2}^{i_2} \dots z_{j+\ell+1}^{i_\ell} \frac{a_{\ell-1+i_1+i_2+\dots+i_\ell}}{a_n} \right| \\
 &\leq \sum_{s=0}^{\infty} \left| \frac{a_{\ell-1+s}}{a_n} \right| R^s \sum_{\substack{i_1 \dots i_\ell \geq 0, \\ i_1+i_2+\dots+i_\ell=s}} 1 \\
 (2.6) \quad &= \sum_{s=0}^{\infty} \left| \frac{a_{\ell-1+s}}{a_n} \right| R^s \left| \binom{-\ell}{s} \right|.
 \end{aligned}$$

If  $k \geq j$  then all indices  $\ell - 1 + s \geq n$ , so (2.2) gives

$$\begin{aligned}
 \left| \frac{f_{j,n+k}}{a_n} \right| &\leq \sum_{s=0}^{\infty} \rho_{n+1}^{-(\ell-1-n+s)} R^s \left| \binom{-\ell}{s} \right| \\
 &= \rho_{n+1}^{-(\ell-1-n)} (1 - R/\rho_{n+1})^{-\ell}.
 \end{aligned}$$

Then

$$(2.7) \quad (\rho_n \rho_{n+1})^{\frac{k-j}{2}} \left| \frac{f_{j,n+k}}{a_n} \right| \leq \left( \frac{\rho_n}{\rho_{n+1}} \right)^{\frac{k-j}{2}} (1 - R/\rho_{n+1})^{-(n+1+k-j)}.$$

If  $k < j$ , then we split into those indices  $<$  and  $\geq n - 1$ , and use the appropriate inequalities in (2.2):

$$\begin{aligned}
 \left| \frac{f_{j,n+k}}{a_n} \right| &\leq \sum_{s=0}^{j-k-1} \rho_n^{-(\ell-1-n+s)} R^s \left| \binom{-\ell}{s} \right| + \sum_{s=j-k}^{\infty} \rho_{n+1}^{-(\ell-1-n+s)} R^s \left| \binom{-\ell}{s} \right| \\
 &\leq \sum_{s=0}^{j-k-1} \rho_n^{-(k-j+s)} R^s \left| \binom{-\ell}{s} \right| + \sum_{s=j-k}^{\infty} \rho_n^{-(k-j+s)} R^s \left| \binom{-\ell}{s} \right| \\
 &= \rho_n^{-(k-j)} (1 - R/\rho_n)^{-\ell}.
 \end{aligned}$$

Then

$$(\rho_n \rho_{n+1})^{\frac{k-j}{2}} \left| \frac{f_{j,n+k}}{a_n} \right| \leq \left( \frac{\rho_n}{\rho_{n+1}} \right)^{\frac{|k-j|}{2}} (1 - R/\rho_n)^{-(n+1+k-j)}.$$

Since  $\rho_n \leq \rho_{n+1}$ , this and (2.7) give (2.4).

(b) From (2.6) with  $\ell = n + 1$ ,

$$\begin{aligned} \left| \frac{f_{j,n+j}}{a_n} - 1 \right| &\leq \sum_{s=1}^{\infty} \left| \frac{a_{n+s}}{a_n} \right| R^s \left| \binom{-n-1}{s} \right| \\ &\leq \sum_{s=1}^{\infty} (R/\rho_{n+1})^s \left| \binom{-n-1}{s} \right| = (1 - R/\rho_{n+1})^{-(n+1)} - 1. \end{aligned}$$

■

In estimating the denominators, we need the notion of diagonal dominance: a matrix

$$B = [b_{jk}]_{1 \leq j, k \leq n}$$

is called **diagonally dominant** if for all  $1 \leq j \leq n$ , we have

$$|b_{jj}| > \sum_{k=1, k \neq j}^n |b_{jk}|.$$

We shall use the basic fact that a diagonally dominant matrix has non-zero determinant [17, p. 373].

**Lemma 2.3**

Assume that for some  $\varepsilon \in (0, \frac{1}{2})$ , infinite sequence of integers  $\mathcal{S}$ , and  $n = j_i, i \in \mathcal{S}$ ,

$$(2.8) \quad \frac{\rho_n}{\rho_{n+1}} \leq \frac{1}{9} (1 - \varepsilon);$$

and

$$(2.9) \quad \lim_{n=j_i, i \in \mathcal{S}} \frac{\rho_n}{n} = \infty.$$

Then for any  $R > 0$ , and all  $\Lambda_n \subset \overline{B_R}$ , we have for large enough  $n = j_i, i \in \mathcal{S}$ ,

$$(2.10) \quad \inf_{|z| \leq R} |q_n(\Lambda_n, z)| > 0.$$

**Proof**

Let us assume that the integers  $\{j_i\} = \{j_i(f)\}$  and  $\{\rho_n\}$  are chosen as above. Assume that  $n = j_i$  for some  $i$ . We use (1.9):

$$q_n(\Lambda_n, z) = \det[\mathbf{c} B]$$

where  $\mathbf{c}$  is an  $(n+1) \times 1$  column vector, and  $B$  is an  $(n+1) \times n$  matrix:

$$\mathbf{c} = \left[ \prod_{\ell=1}^{j-1} (z - z_\ell) \right]_{1 \leq j \leq n+1} ; B = [f_{j-1, n+k-1}]_{\substack{1 \leq j \leq n+1; \\ 2 \leq k \leq n+1}}.$$

We divide the 2nd, 3rd, ...,  $(n+1)$ st column in  $[\mathbf{c} \ B]$  by  $a_n$ , and multiply the  $j$ th row by  $(\rho_{n+1}\rho_n)^{-j/2}$  and  $k$ th column by  $(\rho_{n+1}\rho_n)^{k/2}$  for all  $j, k$ . Then we obtain

$$q_n(\Lambda_n, z) a_n^{-n} = \det [\hat{\mathbf{c}} \ \hat{B}],$$

where

$$\hat{\mathbf{c}} = \left[ (\rho_{n+1}\rho_n)^{(1-j)/2} \prod_{\ell=1}^{j-1} (z - z_\ell) \right]_{1 \leq j \leq n+1} ; \hat{B} = \left[ (\rho_{n+1}\rho_n)^{(k-j)/2} \frac{f_{j-1, n+k-1}}{a_n} \right]_{\substack{1 \leq j \leq n+1; \\ 2 \leq k \leq n+1}}.$$

Let  $R > 0$ ,  $|z| \leq R$ , and all  $|z_j| \leq R$ . We now show that this last matrix  $[\hat{\mathbf{c}} \ \hat{B}]$  is diagonally dominant for  $n = j_i$  and  $i$  large enough. Consider the  $j$ th row. If  $j = 1$ , its diagonal element is 1. For  $j \geq 2$ , the diagonal element is

$$(2.11) \quad \frac{f_{j-1, n+j-1}}{a_n} = 1 + \varepsilon_{j, n},$$

where by the previous lemma, and by (2.9),

$$(2.12) \quad |\varepsilon_{j, n}| \leq (1 - R/\rho_{n+1})^{-n} - 1 \rightarrow 0 \text{ as } n \rightarrow \infty,$$

uniformly in  $j$ . Now consider the sum of the absolute values of the non-diagonal elements in the  $j$ th row of  $[\hat{\mathbf{c}} \ \hat{B}]$ , namely

$$\tau_j := (\rho_{n+1}\rho_n)^{(1-j)/2} \left| \prod_{\ell=1}^{j-1} (z - z_\ell) \right| + \left( \sum_{k=2}^{j-1} + \sum_{k=j+1}^{n+1} \right) (\rho_{n+1}\rho_n)^{(k-j)/2} \left| \frac{f_{j-1, n+k-1}}{a_n} \right|.$$

Of course, if  $j = 1$ , the first term and first sum are omitted. Using that  $|z| \leq R$ , and all  $|z_k| \leq R$ , and (2.4), we continue this as

$$\begin{aligned}
\tau_j &\leq \left( \frac{2R}{(\rho_{n+1}\rho_n)^{1/2}} \right)^{j-1} + \left( \sum_{k=2}^{j-1} + \sum_{k=j+1}^{n+1} \right) \left( \frac{\rho_n}{\rho_{n+1}} \right)^{\frac{|k-j|}{2}} (1 - R/\rho_n)^{-(n+1+k-j)} \\
&\leq \frac{2R}{(\rho_{n+1}\rho_n)^{1/2}} + (1 - R/\rho_n)^{-2n-2} 2 \sum_{\ell=1}^{\infty} \left( \frac{\rho_n}{\rho_{n+1}} \right)^{\ell/2} \\
&= \frac{2R}{(\rho_{n+1}\rho_n)^{1/2}} + (1 - R/\rho_n)^{-2n-2} \frac{2 \left( \frac{\rho_n}{\rho_{n+1}} \right)^{1/2}}{1 - \left( \frac{\rho_n}{\rho_{n+1}} \right)^{1/2}} \\
&< o(1) + (1 + o(1)) \frac{\frac{2}{3}(1 - \varepsilon)^{1/2}}{1 - \frac{1}{3}(1 - \varepsilon)^{1/2}} < (1 + o(1))(1 - \varepsilon)^{1/2},
\end{aligned}$$

by (2.8) and (2.9). In view of (2.11), (2.12), we have diagonal dominance of  $[\hat{\mathbf{c}} \hat{B}]$ , and then (2.10) follows. ■

#### Proof of Theorem 1.4

Now for  $n = j_i$ , for some  $i \geq 1$ , we have

$$|a_n| = K / \prod_{\ell=1}^n \rho_\ell \geq K / \rho_n^n$$

$$\Rightarrow \liminf_{n \rightarrow \infty, n=j_i} \rho_n^{1/n} \geq \liminf_{n \rightarrow \infty, n=j_i} K^{1/n^2} / |a_n|^{1/n^2} \geq 3 / (1 - \varepsilon)^{1/2},$$

for some  $\varepsilon > 0$ , by (1.7). Thus  $\rho_{j_i}$  grows roughly at least as fast as  $(3(1 - \varepsilon)^{1/2})^{j_i}$ . Next, for  $n = j_i$ ,

$$\begin{aligned}
|a_n| &= \frac{K}{\rho_1^n} \prod_{\ell=1}^n \frac{\rho_1}{\rho_\ell} = \frac{K}{\rho_1^n} \prod_{\ell=2}^n \left( \frac{\rho_1}{\rho_2} \frac{\rho_2}{\rho_3} \frac{\rho_3}{\rho_4} \dots \frac{\rho_{\ell-1}}{\rho_\ell} \right) \\
&= \frac{K}{\rho_1^n} \left( \frac{\rho_1}{\rho_2} \right)^{n-1} \left( \frac{\rho_2}{\rho_3} \right)^{n-2} \left( \frac{\rho_3}{\rho_4} \right)^{n-3} \dots \frac{\rho_{n-1}}{\rho_n}.
\end{aligned}$$

Then from (1.7), for some  $\varepsilon > 0$ ,

$$\begin{aligned} \frac{(1-\varepsilon)^{1/2}}{3} &\geq \limsup_{n \rightarrow \infty} |a_n|^{1/n^2} \\ &\geq \liminf_{n \rightarrow \infty} \left( \frac{\rho_n}{\rho_{n+1}} \right)^{[(n-1)+(n-2)+(n-3)\dots+1]/n^2} \\ &= \liminf_{n \rightarrow \infty} \left( \frac{\rho_n}{\rho_{n+1}} \right)^{1/2}. \end{aligned}$$

So for some infinite sequence of integers  $\mathcal{S}$ ,

$$\frac{\rho_n}{\rho_{n+1}} \leq \frac{1-\varepsilon}{9}, \quad n \in \mathcal{S}.$$

So we have (2.8). But since  $\rho_{n+1} = \rho_n$  unless  $n = j_i$  and  $n+1 = j_i+1$  for some  $i$ , so, as above,  $\rho_n = \rho_{j_i}$  grows at least as fast as  $(3(1-\varepsilon))^{j_i}$ . Then (2.9) also follows.

Now let  $S > R > 0$ . It follows from the previous lemma that for  $z \in B_S$  and  $\Lambda_n \subset B_R$ , we have for large enough  $n \in \mathcal{S}$ , that  $|q_n(\Lambda_n, z)| > 0$ , so  $R_n(\Lambda_n, z)$  has no poles in  $B_S$ . Then the uniform convergence in compact subsets of  $B_S$  follows easily from the contour integral error formula for multipoint Padé approximation [2]. ■

We turn to the proof of Theorem 1.1, and first introduce some notation. Recall that  $\mathcal{A}$  denotes the space of entire functions, with metric defined by (1.1). Given  $R > 0$ ,  $n \geq 1$ , we let

$$\mathcal{B}_{n,R} = \{f \in \mathcal{A} : q_n(\Lambda_n, z) \text{ has full degree } n \text{ and no zeros in } \overline{B_R} \text{ whenever } \Lambda_n \subset \overline{B_R}\}.$$

Also, let

$$\mathcal{C}_{n,R} = \bigcup_{j=n}^{\infty} \mathcal{B}_{j,R}.$$

#### Lemma 2.4

For each  $n$  and  $R > 0$ ,  $\mathcal{C}_{n,R}$  is open and dense in  $\mathcal{A}$ .

#### Proof

We first show that each  $\mathcal{B}_{n,R}$  is open, and then the openness of  $\mathcal{C}_{n,R}$  follows. Fix an  $n$  and  $f \in \mathcal{B}_{n,R}$ . Since we need to indicate dependence of the multipoint Padé denominators on  $f$ , we use the notation  $q_n(f, \Lambda_n, z)$  in this proof only. By compactness, and the continuity of  $q_n(f, \Lambda_n, z)$  in  $\Lambda_n$  as long as it has full degree (as follows from (1.8), (1.9)), we see that

$$\min \{|q_n(f, \Lambda_n, z)| : z \in \overline{B_R}, \Lambda_n \subset \overline{B_R}\} > 0.$$

Moreover, by our hypothesis that the denominators have full degree, all their leading coefficients are non-zero, and then also from (1.9).

$$\min \left\{ \left| \det [f_{j-1, n+k}]_{1 \leq j, k \leq n} \right| : \Lambda_n \subset \overline{B_R} \right\} > 0.$$

If we consider entire  $g$  with  $d^*(f, g)$  small enough, then the Maclaurin series coefficients of  $g$  will be as close to those of  $f$  as we please, and consequently all finite differences  $g_{j-1, n+k}$  will be close to the corresponding differences for  $f$ . Then we can ensure that also

$$\min \left\{ \left| \det [g_{j-1, n+k}]_{1 \leq j, k \leq n} \right| : \Lambda_n \subset \overline{B_R} \right\} > 0.$$

By the same token, considering the formula (1.9), we can also ensure that

$$\min \left\{ |q_n(g, \Lambda_n, z)| : z \in \overline{B_R}, \Lambda_n \subset \overline{B_R} \right\} > 0.$$

Thus also  $g \in \mathcal{B}_{n, R}$ . So each  $\mathcal{B}_{n, R}$  is open, and hence each  $\mathcal{C}_{n, R}$  is open.

The denseness is somewhat more difficult. Fix now some  $f \in \mathcal{A}$ . We shall construct  $g \in \mathcal{C}_{n, R}$  with  $d^*(f, g)$  as small as we please. Let us assume that the integers  $\{j_i\} = \{j_i(f)\}$  and  $\{\rho_n\} = \{\rho_n(f)\}$  are chosen as above. Choose now some large positive integer  $i_0$ . Note that by choice of our  $\{\rho_j\}$  above,

$$|a_n| \leq K / \prod_{j=1}^n \rho_j(f) \text{ for } n \leq j_{i_0}(f)$$

and we have equality when  $n = j_k$ , some  $1 \leq k \leq i_0$ . Now define  $\hat{\rho}_j$  for  $j > j_{i_0}$  by

$$\hat{\rho}_j = 10^{j-j_{i_0}} \rho_{j_{i_0}}.$$

Also define for  $n > j_{i_0}$ ,

$$\hat{a}_n = a_{j_{i_0}} / \prod_{j=j_{i_0}+1}^n \hat{\rho}_j = a_{j_{i_0}} \rho_{j_{i_0}}^{-(n-j_{i_0})} 10^{-(n-j_{i_0})(n-j_{i_0}+1)/2}$$

and

$$g(z) = \sum_{j=0}^{j_{i_0}} a_j z^j + \sum_{j=j_{i_0}+1}^{\infty} \hat{a}_j z^j.$$

Then  $g$  is entire (of order 0), and

$$d^*(f, g) \leq \sup \left\{ \left| a_n - a_{j_{i_0}} \rho_{j_{i_0}}^{-(n-j_{i_0})} 10^{-(n-j_{i_0})(n-j_{i_0}+1)/2} \right|^{1/n} : n > j_{i_0} \right\}.$$

Straightforward estimation shows that by choosing  $i_0$  large enough, this can be made as small as we please. Next, as  $g$  and  $f$  have the same series coefficients up to the coefficient of  $z^{j_{i_0}}$ , we see that

$$j_k(g) = j_k(f), k \leq i_0 \text{ and } \rho_n(g) = \rho_n(f), n \leq j_{i_0}.$$

Next, if  $n > m \geq j_{i_0}$ , we see that

$$\begin{aligned} \left( \frac{|\hat{a}_n|}{|\hat{a}_m|} \right)^{1/(n-m)} &= \left( \prod_{j=m+1}^n \left( \rho_{j_{i_0}}^{-1} 10^{-(j-j_{i_0})} \right) \right)^{1/(n-m)} \\ &= \rho_{j_{i_0}}^{-1} 10^{j_{i_0}} \left( 10^{-n(n+1)/2+m(m+1)/2} \right)^{1/(n-m)} \\ &= \rho_{j_{i_0}}^{-1} 10^{j_{i_0}} 10^{-(n-m)(n+m+1)/(2(n-m))} \\ &= \rho_{j_{i_0}}^{-1} 10^{j_{i_0} - (n+m+1)/2} \end{aligned}$$

which is maximal for a given  $m$  and  $n > m$ , iff  $n = m + 1$ . It follows easily that

$$j_{i_0+k}(g) = j_{i_0}(f) + k, k \geq 1,$$

and for  $n > j_{i_0}$ ,

$$\rho_n(g) = \hat{\rho}_n = 10^{n-j_{i_0}} \rho_{j_{i_0}}.$$

Then for  $n \geq i_0$ ,

$$\frac{\rho_{n+1}(g)}{\rho_n(g)} = 10$$

and

$$\lim_{n \rightarrow \infty} \rho_n(g)^{1/n} = 10.$$

It then follows from Lemma 2.3, that for large enough  $n$  and all  $\Lambda_n \subset \overline{B_R}$ ,  $q_n(g, \Lambda_n, z)$  has no zeros in  $\overline{B_R}$ . Thus  $g \in \mathcal{B}_{n,R}$  for all large enough  $n$ , and in particular,  $g \in \mathcal{C}_{n,R}$ , while  $g$  may be made as close to the given  $f$  as we please, by choosing  $i_0$  large enough. As  $f \in \mathcal{A}$  is arbitrary, so  $\mathcal{C}_{n,R}$  is dense. ■

### Proof of Theorem 1.1

Let

$$\mathcal{C} = \bigcap_{\ell=1}^{\infty} \bigcap_{n=1}^{\infty} \mathcal{C}_{n,\ell}.$$

and

$$\mathcal{E} = \mathcal{A} \setminus \mathcal{C} = \bigcup_{\ell=1}^{\infty} \bigcup_{n=1}^{\infty} \mathcal{A} \setminus \mathcal{C}_{n,\ell}.$$

Here since  $\mathcal{C}_{n,\ell}$  is open and dense,  $\mathcal{E} = \mathcal{A} \setminus \mathcal{C}$  is a countable union of closed nowhere dense sets, and is an  $F_\sigma$  set. Next if  $f \in \mathcal{A} \setminus \mathcal{E}$ , then  $f \in \mathcal{C}$ , so  $f \in \mathcal{C}_{n,\ell}$  for all  $n, \ell$ . Then we can choose an increasing sequence

of integers  $\{n_\ell\}_{\ell \geq 1}$  such that  $f \in \mathcal{B}_{n_\ell, \ell}$  for  $\ell \geq 1$ . Then  $q_{n_\ell}(\Lambda_{n_\ell}, z)$  has full degree  $n$  and no zeros in  $\overline{B_\ell}$  whenever  $\Lambda_{n_\ell} \subset \overline{B_\ell}$ . This gives the desired uniform convergence of  $\{R_{n_\ell}(\Lambda_{n_\ell}, \cdot)\}_{\ell \geq 1}$  whenever  $\Lambda_{n_\ell} \subset \overline{B_R}$ , some  $R > 0$ . ■

### 3. PROOF OF THEOREM 1.2

We proceed partly as in [20]. Let  $a_j \neq 0, j \geq 1$ , and

$$(3.1) \quad \sigma_j = a_{j-1}a_{j+1}/a_j^2, \quad j \geq 1.$$

For integers  $t \geq -n + 1$ , we let

$$(3.2) \quad r_{n,t} = \frac{a_{n+t}}{a_n} \left( \frac{a_n}{a_{n+1}} \sigma_n^{1/2} \right)^t.$$

#### Lemma 3.1

(a)

$$(3.3) \quad r_{n,t} = \sigma_n^{|t|/2} \prod_{\ell=1}^{|t|-1} \sigma_{n+\ell \text{sign}(t)}^{|t|-\ell}.$$

(b) Assume that for  $j \geq J$ ,

$$(3.4) \quad |\sigma_j| \leq \chi < 1.$$

Then for  $n \geq 1$  and  $t \geq -n - 1$  such that  $\min\{n, n + t + 1\} \geq J$ ,

$$(3.5) \quad |r_{n,t}| \leq \chi^{t^2/2}.$$

(c) If (3.5) holds for  $j \geq J$ , then for some  $C_0 = C_0(J)$ , and all  $n \geq J$ , and  $t = 0, \pm 1, \pm 2, \dots$  such that  $n + t > 0$ ,

$$(3.6) \quad |r_{n,t}| \leq C_0 \chi^{t^2/2}.$$

#### Proof

(a) If  $t > 0$ , we use

$$\frac{a_{j+1}}{a_j} = \sigma_j \frac{a_j}{a_{j-1}},$$



so

$$\begin{aligned}
\frac{a_{n+t}}{a_n} &= \prod_{k=0}^{t-1} \frac{a_{n+k+1}}{a_{n+k}} \\
&= \prod_{k=0}^{t-1} \left( \sigma_{n+k} \sigma_{n+k-1} \cdots \sigma_{n+1} \frac{a_{n+1}}{a_n} \right) \\
(3.7) \quad &= \left( \frac{a_{n+1}}{a_n} \right)^t \sigma_{n+1}^{t-1} \sigma_{n+2}^{t-2} \cdots \sigma_{n+t-1}.
\end{aligned}$$

If  $t < 0$ , we use

$$\frac{a_{j-1}}{a_j} = \sigma_j \frac{a_j}{a_{j+1}},$$

so

$$\begin{aligned}
\frac{a_{n+t}}{a_n} &= \prod_{k=1}^{|t|} \frac{a_{n-k}}{a_{n-k+1}} \\
&= \prod_{k=1}^{|t|} \left( \sigma_{n-k+1} \sigma_{n-k+2} \cdots \sigma_n \frac{a_n}{a_{n+1}} \right) \\
&= \left( \frac{a_n}{a_{n+1}} \right)^{|t|} \sigma_n^{|t|} \sigma_{n-1}^{|t|-1} \cdots \sigma_{n+1-|t|}.
\end{aligned}$$

This and (3.7) easily give the identity (3.3).

(b) This follows directly from (a), as  $n + \ell \text{sign}(t) \geq \min\{n, n+t-1\} \geq J$  in the product, so we can apply the bound (3.4).

(c) Assume

$$C_1 = \sup_{j \geq 1} |\sigma_j|.$$

This also follows directly from (a). Indeed, using (3.4),

$$|r_{n,t}| \leq \chi^{|t|/2} \left( \prod_{\substack{1 \leq \ell \leq |t|-1; \\ n + \ell \text{sign}(t) \geq J}} \chi^{|t|-\ell} \right) \prod_{\substack{1 \leq \ell \leq |t|-1; \\ n + \ell \text{sign}(t) < J}} C_1^{|t|-\ell}$$

which easily gives the result, as there are  $O(J^2)$  factors of  $C_1$  arising from  $\ell$  where  $n + \ell \text{sign}(t) < J$ . ■

Next for given  $n$  and  $j \geq 0, k \geq 0$ , let

$$(3.8) \quad b_{j,k} = \frac{f_{j,n+k}}{a_n} \left( \frac{a_n}{a_{n+1}} \sigma_n^{1/2} \right)^{k-j},$$

where  $f_{j,n+k}$  is the divided difference in (1.8).

**Lemma 3.2**

Assume all  $|z_j| \leq R$ . Assume also (3.4) holds for  $j \geq J$ .

(a) There exists  $M$  depending only on  $R, \chi, J$ , such that for  $0 \leq j, k \leq n$ , satisfying

$$(3.9) \quad n + k - j \geq M,$$

we have for some constant  $C_2$  depending only on  $R, \chi, J$ ,

$$(3.10) \quad |b_{j,k}| \leq \chi^{(k-j)^2/2} \{1 + C_2(n+1+k-j)\chi^{n+1+k-j}\}.$$

(b) For  $n \geq M$  and  $0 \leq j \leq n$ ,

$$(3.11) \quad |b_{j,j} - 1| \leq C_2 n \chi^n.$$

(c) We have for some constant  $C_4$  depending only on  $R, \chi, J$ , and all  $0 \leq j, k \leq n$ ,

$$(3.12) \quad |b_{j,k}| \leq C_4 \chi^{(k-j)^2/2}.$$

**Proof**

(a) We may assume that  $M > J$ . Let  $\ell = n + 1 + k - j$ . Using (2.6),

$$\left| \frac{f_{j,n+k}}{a_n} \right| \leq \sum_{s=0}^{\infty} \left| \frac{a_{\ell-1+s}}{a_n} \right| R^s \left| \binom{-\ell}{s} \right|.$$

Then as all  $\ell - 1 + s \geq J$ , we can apply our definitions (3.2), (3.8), and the bound (3.5) and deduce

$$\begin{aligned} |b_{j,k}| &\leq \sum_{s=0}^{\infty} \left( \left| \frac{a_n}{a_{n+1}} \right| \sigma_n^{1/2} \right)^{-s} |r_{n,k-j+s}| R^s \left| \binom{-\ell}{s} \right| \\ &\leq \sum_{s=0}^{\infty} \left( \left| \frac{a_n}{a_{n+1}} \right| \sigma_n^{1/2} \right)^{-s} \chi^{(k-j+s)^2/2} R^s \left| \binom{-\ell}{s} \right| \\ &= \chi^{(k-j)^2/2} \sum_{s=0}^{\infty} \left( \left| \frac{a_{n+1}}{a_n} \right| \sigma_n^{-1/2} \chi^{k-j} R \right)^s \chi^{s^2/2} \left| \binom{-\ell}{s} \right| \\ &\leq \chi^{(k-j)^2/2} \sum_{s=0}^{\infty} \left( \left| \frac{a_{n+1}}{a_n} \right| \sigma_n^{-1/2} \chi^{k-j+1/2} R \right)^s \left| \binom{-\ell}{s} \right|. \end{aligned}$$

Here

$$\begin{aligned}
 & \left| \frac{a_{n+1}}{a_n} \right| \sigma_n^{-1/2} \chi^{k-j+1/2} R \\
 &= \sigma_n \sigma_{n-1} \sigma_{n-2} \dots \sigma_J \left| \frac{a_J}{a_{J-1}} \right| \sigma_n^{-1/2} \chi^{k-j+1/2} R \\
 &\leq \chi^{n+1+k-j-J} \left| \frac{a_J}{a_{J-1}} \right| R \\
 (3.13) \quad &= C_3 \chi^\ell,
 \end{aligned}$$

where  $C_3 = \chi^{-J} \left| \frac{a_J}{a_{J+1}} \right| R$ . So

$$\begin{aligned}
 |b_{j,k}| &\leq \chi^{(k-j)^2/2} \sum_{s=0}^{\infty} (C_3 \chi^\ell)^s \left| \binom{-\ell}{s} \right| \\
 &\leq \chi^{(k-j)^2/2} (1 - C_3 \chi^\ell)^{-\ell},
 \end{aligned}$$

provided  $C_3 \chi^\ell < 1$ . We now use the inequality

$$(3.14) \quad (1-x)^{-\ell} - 1 \leq 2\ell x \text{ for } x \in \left[0, \frac{1}{2\ell}\right], \ell \geq 2.$$

This gives

$$|b_{j,k}| \leq \chi^{(k-j)^2/2} \{1 + 2C_3 \ell \chi^\ell\},$$

provided

$$(3.15) \quad C_3 \chi^\ell \leq \frac{1}{2\ell}.$$

As  $\chi < 1$ , this last inequality holds for  $\ell = n + 1 + k - j \geq M$  and some threshold  $M$ .

(b) Here, proceeding as in (a), we see that for  $n \geq J$ , with  $\ell = n + 1$ ,

$$\begin{aligned}
 \left| \frac{f_{j,n+j}}{a_n} - 1 \right| &\leq \sum_{s=1}^{\infty} \left| \frac{a_{\ell-1+s}}{a_n} \right| R^s \left| \binom{-\ell}{s} \right| \\
 &\leq \sum_{s=1}^{\infty} \left( \left| \frac{a_n}{a_{n+1}} \right| \sigma_n^{1/2} \right)^{-s} |r_{n,s}| R^s \left| \binom{-\ell}{s} \right| \\
 &\leq \sum_{s=1}^{\infty} \left( \left| \frac{a_n}{a_{n+1}} \right| \sigma_n^{1/2} \right)^{-s} \chi^{s^2/2} R^s \left| \binom{-\ell}{s} \right| \\
 &\leq \sum_{s=1}^{\infty} \left( \left| \frac{a_{n+1}}{a_n} \right| \sigma_n^{-1/2} R \chi^{1/2} \right)^s \left| \binom{-\ell}{s} \right| \\
 &\leq (1 - C_3 \chi^n R)^{-n-1} - 1,
 \end{aligned}$$

as above. Using (3.14), we continue this for  $n \geq M$ , and uniformly in  $j$ , as

$$|b_{j,j} - 1| \leq C_2 n \chi^n.$$

(c) As above, we let  $\ell = n + 1 + k - j$ . If  $n + 1 + k - j \geq M$ , we can apply (a). So assume  $\ell = n + 1 + k - j < M$ . Now

$$\left| \binom{-\ell}{s} \right| = \binom{\ell + s - 1}{s} \leq 2^{\max\{\ell + s - 1, 0\}} \leq 2^{\ell + s}.$$

Proceeding as in (a), but using Lemma 3.1(c),

$$\begin{aligned} |b_{j,k}| &\leq \sum_{s=0}^{\infty} \left( \left| \frac{a_n}{a_{n+1}} \right| \sigma_n^{1/2} \right)^{-s} |r_{n,k-j+s}| R^s \left| \binom{-\ell}{s} \right| \\ &\leq C_0 \sum_{s=0}^{\infty} \left( \left| \frac{a_n}{a_{n+1}} \right| \sigma_n^{1/2} \right)^{-s} \chi^{(k-j+s)^2/2} R^s 2^{\ell+s} \\ &\leq C_0 2^\ell \chi^{(k-j)^2/2} \sum_{s=0}^{\infty} \left( 2 \left| \frac{a_{n+1}}{a_n} \right| \sigma_n^{-1/2} \chi^{k-j+1/2} R \right)^s \chi^{(s^2-s)/2}. \end{aligned}$$

As at (3.13), and since  $0 \leq \ell = n + 1 + k - j < M$ ,

$$2 \left| \frac{a_{n+1}}{a_n} \right| \sigma_n^{-1/2} \chi^{k-j+1/2} R \leq 2R \chi^{n+k-j-J+1} \left| \frac{a_J}{a_{J-1}} \right| \leq 2R \chi^{-J} \left| \frac{a_J}{a_{J-1}} \right| = 2C_3,$$

so

$$|b_{j,k}| \leq C_0 2^M \chi^{(k-j)^2/2} \sum_{s=0}^{\infty} (2C_3)^s \chi^{(s^2-s)/2}.$$

Then (3.12) follows. ■

### Lemma 3.3

Assume that (1.4) holds with  $\chi < \rho_0$ , where  $\rho_0 = 0.4559\dots$  is the root of (1.5). For any  $R > 0$ , and  $\Lambda_n \subset \overline{B_R}$ , we have for large enough  $n$ ,

$$(3.16) \quad \inf_{|z| \leq R} |q_n(\Lambda_n, z)| > 0.$$

### Proof

We use (1.9):

$$q_n(\Lambda_n, z) = \det[\mathbf{c} B]$$

where  $\mathbf{c}$  is a column vector, and  $B$  is an  $(n+1) \times n$  matrix:

$$\mathbf{c} = \left[ \prod_{\ell=1}^{j-1} (z - z_\ell) \right]_{1 \leq \ell \leq n+1} ; B = [f_{j-1, n+k-1}]_{\substack{1 \leq j \leq n+1; \\ 2 \leq k \leq n+1}}$$

We divide the 2nd, 3rd, ...,  $(n+1)$ st column by  $a_n$ , and multiply the  $j$ th row by  $\left(\frac{a_n}{a_{n+1}}\sigma_n^{1/2}\right)^{-j}$  and  $k$ th column by  $\left(\frac{a_n}{a_{n+1}}\sigma_n^{1/2}\right)^k$ , for all  $j, k$ . Then we obtain

$$(3.17) \quad q_n(\Lambda_n, z) a_n^{-n} = \det \left[ \hat{\mathbf{c}} \hat{B} \right],$$

where

$$\hat{\mathbf{c}} = \left[ \left( \frac{a_n}{a_{n+1}} \sigma_n^{1/2} \right)^{1-j} \prod_{\ell=1}^{j-1} (z - z_\ell) \right]_{1 \leq j \leq n+1};$$

and after an index change,

$$\hat{B} = \left[ \left( \frac{a_n}{a_{n+1}} \sigma_n^{1/2} \right)^{k-j} \frac{f_{j,n+k}}{a_n} \right]_{\substack{0 \leq j \leq n; \\ 1 \leq k \leq n}} = [b_{j,k}]_{\substack{0 \leq j \leq n; \\ 1 \leq k \leq n}}.$$

$$(3.18)$$

Here we are using the notation (3.8). Let  $R > 0$ ,  $|z| \leq R$ , and all  $|z_j| \leq R$ . We now show that  $[\hat{\mathbf{c}} \hat{B}]$  is diagonally dominant. Consider the  $(j+1)$ st row, where  $0 \leq j \leq n$ . If  $j = 0$ , its diagonal element is 1. For  $j \geq 1$ , the diagonal element is

$$(3.19) \quad b_{j,j} = 1 + \varepsilon_{j,n},$$

where by Lemma 3.2(b),

$$(3.20) \quad |\varepsilon_{j,n}| \leq C_2 n \chi^n \rightarrow 0 \text{ as } n \rightarrow \infty,$$

uniformly in  $j$ . Now consider the sum of the absolute values of the non-diagonal elements in the  $(j+1)$ st row, namely

$$\tau_j = \left( \left| \frac{a_{n+1}}{a_n} \sigma_n^{-1/2} \right| \right)^j \left| \prod_{\ell=1}^j (z - z_\ell) \right| + \left( \sum_{k=1}^{j-1} + \sum_{k=j+1}^n \right) |b_{j,k}|.$$

Of course, if  $j = 1$ , the first term and first sum are omitted. First let us assume that

$$n - j \geq \left\lceil \frac{2 \log n}{|\log \chi|} \right\rceil = \Delta_n,$$

say. Then for large enough  $n$ , we have (3.9) for all terms in the sum, so can then estimate

$$\begin{aligned} \tau_j &\leq \left( \left| \frac{a_{n+1}}{a_n} \sigma_n^{-1/2} \right| 2R \right)^j + \left( \sum_{k=1}^{j-1} + \sum_{k=j+1}^{n+1} \right) \chi^{(k-j)^2/2} \{1 + C_2(n+1+k-j)\chi^{n+1+k-j}\} \\ &\leq o(1) + (1 + o(1)) 2 \sum_{\ell=1}^{\infty} \chi^{\ell^2/2} \leq 1 - \varepsilon, \end{aligned}$$

for some small enough  $\varepsilon$ . Recall that  $\chi < \rho_0$ , where  $\rho_0$  is the root of (1.5). Using (3.19), (3.20), we see then that the  $(j+1, j+1)$  element in the  $(j+1)$ st row has absolute value larger than  $\tau_j$ , as required for diagonal dominance. We still have to handle those terms for which  $n-j \leq \Delta_n$ . Here most of  $\tau_j$  can be estimated as above:

$$\left( \left| \frac{a_{n+1}}{a_n} \sigma_n^{-1/2} \right| 2R \right)^j + \sum_{k=\Delta_n+1}^{n+1} \chi^{(k-j)^2/2} \{1 + C_2(n+1+k-j) \chi^{n+1+k-j}\} \leq 1-\varepsilon.$$

Next, for  $2 \leq k \leq \Delta_n$ , and  $n-j \leq \Delta_n$ , we have  $j-k \geq n-2\Delta_n$ , so

$$\begin{aligned} & \sum_{k=2}^{\Delta_n} |b_{j,k}| \\ & \leq C_3 \sum_{k=2}^{\Delta_n} \chi^{(k-j)^2/2} \\ & \leq C_3 \Delta_n \chi^{(n-2\Delta_n)^2/2} \rightarrow 0, \end{aligned}$$

as  $n \rightarrow \infty$ , by Lemma 3.2(c). Again we obtain diagonal dominance. ■

### Proof of Theorem 1.2

Given any  $r > 0$ , the interpolants  $R_n(\Lambda_n, z)$  have no poles in  $B_r$  for large enough  $n$ . Then as above, the locally uniform convergence follows. ■

## 4. PROOF OF THEOREM 1.3

This case is more delicate than the proof of Theorem 1.2. We have to multiply by a suitable matrix before proving diagonal dominance. Accordingly for  $q \in \mathbb{C}$ , and  $n \geq 1$ , let

$$A_n(q) = \left[ q^{(k-j)^2/2} \right]_{1 \leq j, k \leq n}.$$

The determinant of this matrix can be reduced to that of a Vandermonde matrix by multiplying rows and columns by suitable factors. It is known that (see e.g. [21, p. 326])

$$\det(A_n(q)) = \prod_{j=1}^{n-1} (1 - q^j)^{n-j}.$$

When this matrix is non-singular, its inverse admits uniform bounds on its entries. More precisely, the  $(k, \ell)$  entry in  $A_n(q)^{-1}$  admits the bound

$$(4.1) \quad |(A_n(q)^{-1})_{k\ell}| \leq S |q|^{|\ell-k|/2},$$

where

$$S = 2 \prod_{j=1}^{\infty} \left( \frac{1 + |q|^j}{1 - |q|^j} \right)^2.$$

See [21, Lemma 2.1, pp. 326-327]. For integers  $t \geq -n + 1$ , we define  $r_{n,t}$  by (3.2). We begin with bounds and asymptotics for  $r_{n,t}$ :

**Lemma 4.1**

Let  $L \geq 1$ . We have for  $|t| \leq L$ ,

$$r_{n,t} = q^{t^2/2} (1 + \varepsilon_{n,t})$$

where

$$(4.2) \quad \max_{|t| \leq L} |\varepsilon_{n,t}| \rightarrow 0 \text{ as } n \rightarrow \infty.$$

**Proof**

Since the total number of  $\sigma$  factors in the right-hand side of (3.3) is  $|t|/2 + (|t| - 1) + (|t| - 2) + \dots + 1 = |t|^2/2$ , the assertion follows from our hypothesis that  $\sigma_m \rightarrow q$  as  $m \rightarrow \infty$ . ■

Next for given  $n$  and  $j \geq 0, n + k \geq 0$ , define  $b_{j,k}$  by (3.8).

**Lemma 4.2**

Let  $L \geq 1$  and  $\chi \in (|q|, 1)$ . For  $1 \leq j, k \leq n$ , we can write

$$(4.3) \quad b_{j,k} = q^{(k-j)^2/2} + \delta_{j,k},$$

where

(a) if  $|j - k| \leq L$ ,

$$(4.4) \quad \eta_L^* = \max_{|j-k| \leq L} \left| \delta_{j,k} / q^{(k-j)^2/2} \right| \rightarrow 0 \text{ as } n \rightarrow \infty.$$

(b) for all  $0 \leq j, k \leq n$ ,

$$(4.5) \quad |\delta_{j,k}| \leq C_4 \chi^{(k-j)^2/2},$$

where  $C_4$  is independent of  $n, j, k$  and of  $L$  above.

**Proof**

(a) Let  $\ell = n + 1 + k - j$ , where  $|j - k| \leq L$ . As at (2.6),

$$\begin{aligned} \left| \frac{f_{j,n+k}}{a_n} - \frac{a_{\ell-1}}{a_n} \right| &= \left| \frac{a_{\ell-1}}{a_n} \sum_{i_1 \dots i_{\ell} \geq 0 \text{ with at least one } i_j \geq 1} z_{j+1}^{i_1} z_{j+2}^{i_2} \dots z_{j+\ell+1}^{i_{\ell}} \frac{a_{\ell-1+i_1+i_2+\dots+i_{\ell}}}{a_{\ell-1}} \right| \\ &\leq \left| \frac{a_{n+k-j}}{a_n} \sum_{s=1}^{\infty} R^s \left| \frac{a_{\ell-1+s}}{a_{\ell-1}} \right| \left| \binom{-\ell}{s} \right| \right|. \end{aligned}$$

Proceeding as in Lemma 3.2, and using Lemma 3.1(c),

$$\begin{aligned}
|b_{j,k} - r_{n,k-j}| &\leq |r_{n,k-j}| \sum_{s=1}^{\infty} R^s \left| \frac{a_\ell}{a_{\ell-1}} \sigma_{\ell-1}^{-1/2} \right|^s |r_{\ell-1,s}| \left| \binom{-\ell}{s} \right| \\
&\leq |r_{n,k-j}| \sum_{s=1}^{\infty} R^s \left| \frac{a_\ell}{a_{\ell-1}} \sigma_{\ell-1}^{-1/2} \right|^s C_4 \chi^{s^2/2} \left| \binom{-\ell}{s} \right| \\
&\leq |r_{n,k-j}| C_4 \left\{ \left( 1 - R \left| \frac{a_\ell}{a_{\ell-1}} \sigma_{\ell-1}^{-1/2} \right| \chi^{1/2} \right)^{-\ell} - 1 \right\} \\
&\leq |r_{n,k-j}| C_5 (n+k-j) \chi^{n+k-j} \\
&\leq C_6 |r_{n,k-j}| n \chi^n,
\end{aligned}$$

by (3.14). Then Lemma 4.1 gives

$$\begin{aligned}
|\delta_{j,k}| &= \left| b_{j,k} - q^{(k-j)^2/2} \right| \\
&\leq |b_{j,k} - r_{n,k-j}| + \left| r_{n,k-j} - q^{(k-j)^2/2} \right| \\
&\leq |r_{n,k-j}| \left\{ C_6 n \chi^n + C_7 \max_{|t| \leq L} |\varepsilon_{n,t}| \right\} \\
&\leq \eta_L^* |q|^{(k-j)^2/2},
\end{aligned}$$

where  $\eta_L^* \rightarrow 0$  as  $L \rightarrow \infty$ .

(b) Choose  $J$  such that  $|\sigma_j| \leq \chi$  for  $j \geq J$ . Note that  $J$  is independent of  $L$  in (a). Using Lemma 3.2(c), for all  $0 \leq j, k \leq n$ ,

$$|b_{j,k}| \leq C_4 \chi^{(k-j)^2/2}$$

so

$$|\delta_{j,k}| \leq |b_{j,k}| + |q|^{(k-j)^2/2} \leq (C_4 + 1) \chi^{(k-j)^2/2}.$$

■

### Proof of Theorem 1.3

It suffices to show that given  $R > 0$  and  $\Lambda_n \subset B_R$ ,  $n \geq 1$ , then for large enough  $n$ ,

$$\inf_{z \in B_R} |q_n(\Lambda_n, z)| > 0.$$

We use (3.17), namely

$$(4.6) \quad q_n(\Lambda_n, z) a_n^{-n} = \det \left[ \hat{\mathbf{c}} \hat{B} \right],$$



where  $\left[ \hat{\mathbf{c}} \hat{B} \right]$  is given by (3.18). We partition the column  $\hat{\mathbf{c}}$  and matrix  $\hat{B}$  as follows:

$$q_n(\Lambda_n, z) a_n^{-n} = \det \begin{bmatrix} 1 & \mathbf{f} \\ \mathbf{d} & A_n(q) + \Delta \end{bmatrix}$$

where  $\mathbf{d}$  is an  $n$  by 1 column vector;  $\mathbf{f}$  is an  $n$  by 1 row vector; and  $\Delta$  is an  $n$  by  $n$  matrix with "small" entries. Thus

$$\begin{aligned} \mathbf{d} &= \left[ \left( \frac{a_n}{a_{n+1}} \sigma_n^{1/2} \right)^{-j} \prod_{\ell=1}^j (z - z_\ell) \right]_{1 \leq j \leq n} ; \\ \mathbf{f} &= \left[ \left( \frac{a_n}{a_{n+1}} \sigma_n^{1/2} \right)^k \frac{f_{0,n+k}}{a_n} \right]_{1 \leq k \leq n}^T = [b_{0,k}]_{1 \leq k \leq n} ; \\ \Delta &= \left[ \left( \frac{a_n}{a_{n+1}} \sigma_n^{1/2} \right)^{k-j} \frac{f_{j,n+k}}{a_n} \right]_{1 \leq j, k \leq n} - A_n(q) \\ &= [b_{j,k}]_{1 \leq j, k \leq n} - A_n(q) = [\delta_{j,k}]_{1 \leq j, k \leq n} . \end{aligned}$$

(4.7)

We multiply the determinant above by

$$\det \begin{bmatrix} 1 & -\mathbf{f} A_n(q)^{-1} \\ \mathbf{0} & A_n(q)^{-1} \end{bmatrix} = \det A_n(q)^{-1} .$$

We then see that

$$\begin{aligned} & q_n(\Lambda_n, z) a_n^{-n} (\det A_n(q))^{-1} \\ &= \det \begin{bmatrix} 1 & \mathbf{f} \\ \mathbf{d} & A_n(q) + \Delta \end{bmatrix} \begin{bmatrix} 1 & -\mathbf{f} A_n(q)^{-1} \\ \mathbf{0} & A_n(q)^{-1} \end{bmatrix} \\ &= \det \begin{bmatrix} 1 & \mathbf{0}^T \\ \mathbf{d} & -\mathbf{d} \mathbf{f} A_n(q)^{-1} + I + \Delta A_n(q)^{-1} \end{bmatrix} \\ &= \det [I - \mathbf{d} \mathbf{f} A_n(q)^{-1} + \Delta A_n(q)^{-1}] . \end{aligned}$$

(4.8)

We shall show the matrix in this last determinant is diagonally dominant. First,

$$\begin{aligned} & \mathbf{d} \mathbf{f} A_n(q)^{-1} \\ &= \left[ \left( \frac{a_n}{a_{n+1}} \sigma_n^{1/2} \right)^{-j} \left( \prod_{\ell=1}^j (z - z_\ell) \right) b_{0,k} \right]_{1 \leq j, k \leq n} A_n(q)^{-1} \end{aligned}$$

so the sum of the absolute values of elements in the  $j$ th row of the matrix  $\mathbf{d}fA_n(q)^{-1}$  is, using (4.5) and (4.1),

$$\begin{aligned}
& \left| \left( \frac{a_n}{a_{n+1}} \sigma_n^{1/2} \right)^{-j} \prod_{\ell=1}^j (z - z_\ell) \right| \left| \sum_{\ell=1}^n \sum_{k=1}^n b_{0,k} (A_n(q)^{-1})_{k\ell} \right| \\
& \leq SC_3 \left( 2R \left| \frac{a_{n+1}}{a_n} \right| \right)^j \sum_{\ell=1}^n \sum_{k=1}^n \chi^{k^2/2} |q|^{k-\ell/2} \\
& \leq 2SC_3 \left( 1 - |q|^{1/2} \right)^{-1} \left( \sum_{k=1}^{\infty} \chi^{k^2/2} \right) \left( 2R \left| \frac{a_{n+1}}{a_n} \right| \right)^j \\
& \leq C_4 \left( 2R \left| \frac{a_{n+1}}{a_n} \right| \right)^j.
\end{aligned}$$

(4.9)

Next, the sum of absolute values of elements in the  $j$ th row of the matrix  $\Delta A_n(q)^{-1}$  is bounded above by

$$\begin{aligned}
& \sum_{k=1}^n \sum_{\ell=1}^n \left| \delta_{j,\ell} A_n(q)^{-1}_{\ell,k} \right| \\
& \leq S \sum_{k=1}^n \sum_{\ell=1}^n |\delta_{j,\ell}| |q|^{k-\ell/2} \\
& \leq 2S \left( 1 - |q|^{1/2} \right)^{-1} \sum_{\ell=1}^n |\delta_{j,\ell}| \\
& \leq 2S \left( 1 - |q|^{1/2} \right)^{-1} \left\{ \eta_L^* \sum_{\ell:|j-\ell|\leq L} |q|^{(j-\ell)^2/2} + C_3 \sum_{\ell:|j-\ell|>L} \chi^{(j-\ell)^2/2} \right\} \\
& \leq 2S \left( 1 - |q|^{1/2} \right)^{-1} \left\{ 2\eta_L^* \left( 1 - |q|^{1/2} \right)^{-1} + C_3 \chi^{L^2/2} \right\},
\end{aligned}$$

by (4.2) and (4.5). It is crucial here that  $C_3$  is independent of  $L$ . Choosing  $L$  large enough, and then using that  $\eta_L^* \rightarrow 0$  as  $n \rightarrow \infty$ , we see that this row sum may be made  $< \frac{1}{4}$  for large enough  $n$ . Together with (4.9), this shows that the matrix in the determinant in (4.8) is diagonally dominant, and we are done. ■

#### REFERENCES

- [1] A. I. Aptekarev, M. L. Yattselev, *Padé approximants for functions with branch points - strong asymptotics of Nuttall-Stahl polynomials*, Acta Mathematica, 215(2015), 217–280.

- [2] G.A. Baker, P. Graves-Morris, *Padé Approximants*, 2nd Edition, Cambridge University Press, Cambridge, 1996.
- [3] G.A. Baker, *Some structural properties of two counter-examples to the Baker–Gammel–Wills conjecture*, J. Comput. Appl. Math., 161 (2003), 371 – 391.
- [4] G.A. Baker, *Counter-examples to the Baker-Gammel-Wills conjecture and patchwork convergence*, J. Comput. Appl. Math. 179 (2005), 1–14.
- [5] B. Beckermann, G. Labahn, A. C. Matos, *On rational functions without Froissart doublets*, manuscript.
- [6] V. I. Buslaev, *The Baker-Gammel-Wills Conjecture in the Theory of Padé Approximants*, Math. USSR. Sbornik, 193(2002), 811–823.
- [7] V. I. Buslaev, *Convergence of the Rogers-Ramanujan Continued Fraction*, Math. USSR. Sbornik, 194(2003), 833–856.
- [8] V. I. Buslaev, A. A. Gončar, and S. P. Suetin, *On Convergence of Subsequences of the  $m$ th Row of a Padé Table*, Math. USSR Sbornik, 48(1984), 535–540.
- [9] F. Cala Rodrigues, G. Lopez Lagomasino, *Exact Rates of Convergence of Multipoint Padé Approximants*, Constr. Approx., 14(1988), 259-272.
- [10] T. Claeys and F. Wielonsky, *On Sequences of Rational Interpolants of the Exponential Function with Unbounded Interpolation Points*, J. Approx. Theory, 171(2013), 1-32.
- [11] K.A. Driver, *Simultaneous Rational Approximants for a Pair of Functions with Smooth Maclaurin Series Coefficients*, J. Approximation Theory, 83(1995), 308-329.
- [12] J. Gilewicz, Y. Kryakin, *Froissart doublets in Padé approximation in the case of polynomial noise*, J. Comp. Appl. Math., 153(2003), 235–242.
- [13] A. Gonchar, G. Lopez Lagomasino, *Markov’s Theorem for Multipoint Padé Approximants*, Math. Sbornik, 105(1978), 512-524.
- [14] D.V. Khristoforov, *On the Phenomenon of Spurious Interpolation of Elliptic Functions by Diagonal Padé Approximants*, Mathematical Notes, 87(2010), 564-574.
- [15] R. Kovacheva, *Zeros of Padé error functions for functions with smooth Maclaurin coefficients*, J. Approximation Theory, 83(1995), 371-391.
- [16] Yu. Labych and A. Starovoitov, *Trigonometric Padé Approximants for Functions with regularly decreasing Fourier coefficients*, Mathematics Sbornik, 200(2009), 1051-1074.
- [17] P. Lancaster, M. Tismenetsky, *The Theory of Matrices*, Academic Press, San Diego, 1985.
- [18] Eli Levin and D.S. Lubinsky, *Rows and Diagonals of the Walsh Array for Entire Functions with Smooth Maclaurin Series Coefficients*, Constructive Approximation, 6(1990), 257-286.
- [19] D.S. Lubinsky, *Padé Tables of a Class of Entire Functions*, Proc. Amer. Math. Soc., 94(1985), 399-405.
- [20] D.S. Lubinsky, *Padé Tables of Entire Functions of Very Slow and Smooth Growth*, Constr. Approx., 1(1985), 349-358.
- [21] D.S. Lubinsky, *Padé Tables of Entire Functions of Very Slow and Smooth Growth II*, Constr. Approx., 4(1988), 321-339.
- [22] D.S. Lubinsky, *Distribution of Poles of Diagonal Rational Approximants to Functions of Fast Rational Approximability*, Constructive Approximation, 7(1991), 501-519.

- [23] D.S. Lubinsky, *Spurious Poles in Diagonal Rational Approximation*, (in) Progress in Approximation Theory (eds. A.A. Gončar and E.B. Saff), Springer-Verlag, Berlin, 1992, pp. 191-213.
- [24] D.S. Lubinsky, *Rogers-Ramanujan and the Baker-Gammel-Wills (Padé) Conjecture*, Annals of Mathematics, 157(2003), 847-889.
- [25] D.S. Lubinsky, *Exact Interpolation, Spurious Poles, and Uniform Convergence of Multipoint Padé Approximants*, to appear in Math. Sbornik.
- [26] L.M. Milne-Thomson, *The Calculus of Finite Differences*, Chelsea, New York, 1933.
- [27] E. A. Rakhmanov, *On the Convergence of Padé Approximants in Classes of Holomorphic Functions*, Math. USSR Sbornik, 40(1981), 149–155.
- [28] . N. Rusak and A. P. Starovoitov, *Padé approximants for entire functions with regularly decreasing Taylor coefficients*, Mathematics Sbornik 193(9), 63–92 (2002) [Russian Acad. Sci. Sb.Math. 193 (9), 1303–1332 (2002)].
- [29] H. Stahl, *Spurious Poles in Padé Approximation*, J. Comp. Appl. Math., 9(1998), 511-527.
- [30] S.P. Suetin, *Distribution of the Zeros of Padé polynomials and analytic continuation*, Russian Math. Surveys, 70(2015),901-951.
- [31] H. Wallin, *The Convergence of Padé Approximants and the Size of the Power Series Coefficients*, Applicable Anal., 4(1974), 235-251.
- [32] F. Wielonsky, *Riemann-Hilbert Analysis and Uniform Convergence of Rational Interpolants to the Exponential Function*, J. Approx. Theory, 131(2004), 100-148.
- [33] M. Yattselev, *Meromorphic Approximation: Symmetric Contours and Wandering Poles*, manuscript.

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA,  
GA 30332-0160, USA., LUBINSKY@MATH.GATECH.EDU